

**Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño**

F. Zheng and J. Zhu

**Roles of initial ocean surface and subsurface states on successfully predicting 2006–2007 El Niño**

**F. Zheng and J. Zhu**

International Center for Climate and Environment Science (ICCES), Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China

Received: 28 April 2014 – Accepted: 28 May 2014 – Published: 19 June 2014

Correspondence to: F. Zheng (zhengfei@mail.iap.ac.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

The 2006–2007 El Niño event, an unusually weak event, was predicted by most models only after the warming in the eastern Pacific had commenced. In this study, on the basis of an El Niño prediction system, roles of the initial ocean surface and subsurface states on predicting the 2006–2007 El Niño event are investigated to determine conditions favorable for predicting El Niño growth and are isolated in three sets of hindcast experiments. The hindcast is initialized through assimilation of only the sea surface temperature (SST) observations to optimize the initial surface condition (Assim\_SST), only the sea level (SL) data to update the initial subsurface state (Assim\_SL), or both the SST and SL data (Assim\_SST + SL). Results highlight that the hindcasts with three different initial states all can successfully predict the 2006–2007 El Niño event one year in advance and that the Assim\_SST + SL hindcast performs best. A comparison between the various sets of hindcast results further demonstrates that successful prediction is more significantly affected by the initial subsurface state than by the initial surface condition. The accurate initial surface state can easier trigger the prediction of the 2006–2007 El Niño, whereas a more reasonable initial subsurface state can contribute to improving the prediction in the growth of the warm event.

## 1 Introduction

El Niño–Southern Oscillation (ENSO), which is one of the most striking interannual variabilities in the tropical Pacific Ocean, has been studied for several decades. Understanding the changes in its characteristics remains an important issue for worldwide environmental and socioeconomic interests (McPhaden et al., 2006; Ashok and Yamagata, 2009). During the past two to three decades, ENSO forecasts have made remarkable progress, reaching a stage in which skillful predictions can be made 6–12 months in advance. Many climate models from operational centers have routinely been used to make ENSO predictions in real time (Latif et al., 1998; Kirtman et al., 2002),

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

although the skill of sea surface temperature (SST) forecasts in the equatorial Pacific is strongly model dependent and widely divergent across various prediction systems (Jin et al., 2008; Barnston et al., 2012). To improve the prediction skills, intensive effort has been made in both model improvements and better constraint of initial conditions.

5 The latter is particularly important in the ocean because the memory for ENSO resides in the ocean (e.g., Neelin et al., 1998), and the importance of ocean data in making ENSO predictions has been demonstrated in a number of studies (e.g., Ji and Leetmaa, 1997; Ji et al., 2000; Alves et al., 2004; Keenlyside et al., 2005; Zheng et al., 2006; 2007; Zheng and Zhu, 2008; Yang et al., 2010; Zhu et al., 2012).

10 However, as demonstrated by Barnston et al. (2012), an apparent retrogression in skill exists for ENSO predictions in the 2002–2011 study period, compared with those in the 1981–2010 period, and the real-time ENSO prediction skill of models in the past decade is significantly lower than that of the less-advanced models of the 1980s and 1990s. For example, the nine-year sliding correlation for the hindcasts during the 15 1981–2010 period has an average score of 0.65 at a six-month lead time, but the prediction skill of the correlation decreases to 0.42 for the 2002–2011 period (Barnston et al., 2012). Specifically for the 2006–2007 El Niño event, which was unusually weak (Hackert et al., 2007; McPhaden, 2008; Yang et al., 2010), most seasonal climate forecasts from operational centers only predicted the event after the warming had already 20 become apparent and was basinwide (McPhaden, 2008). The composite forecast for December 2006 from the International Research Institute for Climate Prediction (IRI) indicates that the warm anomaly was under-predicted by more than 0.5 °C. (A review of forecast skills is available on the IRI Web site at <http://iri.columbia.edu>.)

25 The limited predictability may be attributed to factors such as errors in the oceanic initial conditions, state-dependent stochastic forcing, or model errors (Palmer, 2001). In this study, we investigated possible improvements in predicting this warm event by focusing on the role of the oceanic initial states. On the basis of the ensemble Kalman filter (EnKF) algorithm (e.g., Evensen, 2009), the roles of the initial ocean surface and subsurface states on the 2006–2007 El Niño predictions were examined in three sets of

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

retrospective forecast experiments. In the first set, only the SST data was assimilated to provide accurate surface initial conditions, whereas only sea level (SL) data was assimilated to provide the updated information of the subsurface water temperature entrained into the mixed layer in the model. In the third set, both SST and SL data were  
 5 assimilated. The forecast differences in the three sets of the retrospective experiments with initializations by the three separate data assimilation schemes were then examined to isolate the effects of the various initial states on predicting the warm event.

## 2 Model and initialization schemes

In this section, we briefly describe the model components and the designs of three  
 10 initialization schemes on the basis of the EnKF algorithm.

### 2.1 Description of the model components

The ENSO coupled model we used here is an intermediate coupled model (ICM) that was first developed by Keenlyside and Kleeman (2002) and Zhang et al. (2005). Its dynamical component consists of both linear and nonlinear components. The former was  
 15 essentially a McCreary-type (1981) modal model but was extended to include horizontally varying background stratification. Different from the commonly used Zebiak–Cane model (1987), this newly developed ICM considers the realistic vertical structure of the upper ocean, spatially varying stratification, and partial nonlinear effects. For example, when 10 vertical modes are included in the calculation of dynamic ocean fields, along  
 20 with a parameterization of the local Ekman-driven upwelling, this model yields realistic simulations of the mean equatorial circulation and its variability in the tropical Pacific (Keenlyside and Kleeman, 2002).

An SST anomaly model is embedded within this dynamical framework, and the governing equation describes the evolution of the mixed-layer temperature anomalies.  
 25 As demonstrated by Zhang et al. (2005), having a realistic parameterization for the

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



temperature of the subsurface water entrained into the mixed-layer ( $T_e$ ) is crucial to the performance of SST simulations in the equatorial Pacific. An empirical  $T_e$  model was constructed from historical data and was demonstrated to be effective for improving the SST simulations. The ocean model was coupled with a statistical atmospheric model that specifically relates wind stress ( $\tau$ ) to SST anomaly fields. All coupled model components exchanged simulated anomaly fields. Information concerning the interactions between the atmosphere ( $\tau$ ) and the ocean SST was exchanged once daily.

### 2.2 Initialization schemes

An EnKF data assimilation system for initializing the ICM was first developed by Zheng et al. (2006) and Zheng et al. (2007). In the system, the EnKF is implemented by using an ensemble square-root filter algorithm with no perturbation of observations (e.g., Evensen, 2004). This EnKF data assimilation scheme was further improved by using a balanced, multivariate model-error approach (Zheng and Zhu, 2008) and was upgraded to use mean preserving transformations in the square-root scheme (Sakov and Oke, 2008; Evensen, 2009). The observed interannual SST and SL anomalies can then be consistently assimilated into the coupled model to provide an accurate initial condition for real-time SST prediction (Zheng and Zhu, 2010). In this study, we performed three initialization schemes to isolate the effects of the various initial states on ENSO prediction (Table 1). The first scheme, which is known as the Assim\_SST, assimilates only the observed SST anomalies into the ICM during the initialization process. In the second scheme, referred to as the Assim\_SL, the model states of the ICM are periodically updated by assimilating only the SL anomaly data by using EnKF. Moreover, both the SST and SL observations are assimilated into the coupled model to initialize the ICM for the ENSO forecast by using the Assim\_SST + SL scheme.

As previously mentioned, the ICM is formed by coupling the ocean model to a simple statistical atmospheric model for interannual wind stress anomalies ( $\tau$ ), and the atmospheric state is a slave only to the modeled SST. Therefore, assimilation of SST observations can provide an accurate initial surface thermal state of ICM and supply a better

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



estimation of the initial atmospheric state. Thus the Assim\_SST scheme in this study represents a better initialization scheme of the surface model states of ICM (Zheng et al., 2006). As demonstrated in Zhang et al. (2005), a distinguished feature added to the dynamical ocean model of ICM for a better performance of SST simulation in the equatorial Pacific is an empirical parameterization for  $T_e$ . Moreover,  $T_e$  can be directly estimated from the modeled SL anomalies by a singular value decomposition (SVD) model developed from the historical relationship between the interannual variations in  $T_e$  and SL. Thus, assimilation of SL data can improve the accuracy of the subsurface state of ICM (Zheng et al., 2007), and the Assim\_SL scheme represents a better initialization of the subsurface model states of ICM. The Assim\_SST + SL scheme is the best initialization scheme of the three different schemes because it can perfect both the surface and subsurface initial states for ICM to provide an ENSO forecast (Zheng and Zhu, 2008). Moreover, to further quantify the relative effects of the accuracy of the surface and subsurface initial states on the 2006–2007 El Niño predictions, the forecast differences between the two single-observational initialization schemes, Assim\_SST and Assim\_SL, and the double-observational initialization scheme, Assim\_SST + SL, were examined to illustrate the relative contributions of the various initial states to ENSO prediction.

### 2.3 Datasets

The observed monthly SST anomaly fields from the extended reconstructed SST (ERSSTv3b; Smith et al., 2008) were assimilated into the coupled model once a month and were also used to verify the model predictions. The monthly-averaged altimeter products were produced by Ssalto/Duacs and were distributed by Aviso, with support from Cnes (<http://www.aviso.oceanobs.com/duacs/>).

### 3 Isolating the impact of oceanic initial conditions on predicting the 2006–2007 El Niño

We initialized the coupled forecasts of the 2006–2007 El Niño event with the three oceanic analyses on 1 January 2006, and all forecasts had the same atmospheric initial condition. Thus, the outcome depended on the effectiveness of the oceanic initial condition for predicting the 2006–2007 El Niño. The initial conditions of anomalous SST and SL from the Assim\_SST, Assim\_SL, and Assim\_SST + SL analysis results are compared with the observations in the top and middle rows in Fig. 1. As expected, for both the amplitude and spatial pattern of the observed SST, the Assim\_SST scheme had a better analysis result than the Assim\_SL scheme. However, the Assim\_SL scheme had a much more similar assimilation result to the observed SL anomaly field than the Assim\_SST scheme. Overall, the Assim\_SST + SL scheme had a more reasonable analysis result in both the SST and SL and can provide a more dynamically consistent initial condition for both the surface and subsurface states (Zheng and Zhu, 2008).

The SST hindcasts are also compared with observations in Fig. 1. After initialization, all three hindcasts developed a warm anomaly event during the entire 12 months of the forecast. All of the hindcast experiments with the three different initial conditions successfully predicted the 2006–2007 El Niño event before 12 months, and the hindcast initialized by the Assim\_SST + SL scheme performed best. The Assim\_SST + SL hindcast exhibited more realistic evolution during the developing and maturing stages of this El Niño and had the earliest and largest warming of the three experiments, with the first warming in June 2006 and a peak anomaly in December 2006. Moreover, this hindcast had a similar magnitude to the observed warming. The hindcast initialized from the Assim\_SL analysis predicted a slightly weaker warming development of the event. Its peak anomaly was approximately 1.3 °C, weaker than that in the Assim\_SST + SL forecast and in the observed SST. With the Assim\_SST analysis, the warm anomaly developed much slower than the prediction initialized by the Assim\_SL

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



analysis or the Assim\_SST + SL analysis, and the peak anomaly of the Assim\_SST hindcast was the weakest among the three hindcast experiments.

To illustrate the effect of ocean assimilation performance on the prediction skill for the 2006–2007 El Niño event, the differences between the three forecasts are shown in Figs. 2 and 3. In terms of the observed development of the warm event, the prediction was best when assimilating both the SST and SL observations into the coupled model, Assim\_SST + SL. Therefore, the forecast differences between the two single-observational initialization schemes, Assim\_SST and Assim\_SL, with the double-observational initialization scheme, Assim\_SST + SL, can help to isolate the relative effects of the accuracy of the surface and subsurface initial states on the 2006–2007 El Niño predictions.

Figure 2 shows the initial and forecast differences estimated between the Assim\_SST + SL and Assim\_SL forecasts for SST, zonal wind stress ( $T_{\text{aux}}$ ), SL, and  $T_e$ , illustrating the effects of the initial surface states, SST and wind stress, on predicting the 2006–2007 El Niño event. For the initial period, a more reasonable zonal SST gradient was captured by assimilating the SST observational data into the model, in which the SST is anomalously warm west of the dateline and cold to the east (Fig. 2c). This type of zonal SST gradient is convenient for generating stronger westerly winds over the equatorial Pacific (Fig. 2d) at the initial time, which results in the triggering of a considerable El Niño event. As indicated by the SL anomalies (Fig. 2b), the strong westerly wind stress anomalies in the lower atmosphere initiated an oceanic downwelling Kelvin wave, which has been alluded to as a facilitator of El Niño (Picaut et al., 2002; Hackert et al., 2007). Forced by the downwelling Kelvin wave, warm water accumulated in the western Pacific tended to sink and began to propagate eastward, resulting in a net warming at the subsurface layers in the eastern equatorial Pacific during the first half of 2006 (Fig. 2b). When the subsurface warm water reached the eastern tropical Pacific and warmed the surface water through vertical mixing in summer 2006, the warm SST activated the local coupled air–sea interactions to produce atmospheric–oceanic anomalies that developed and evolved with continuous warming during fall

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2006. Finally, a 0.4 °C warmer El Niño condition at the end of 2006 was predicted when considering the accuracy of the initial surface ocean–atmosphere condition.

Similarly, the initial and forecasted differences estimated between the Assim\_SST + SL and Assim\_SST forecasts for SL,  $T_e$ , SST, and Taux, are shown in Fig. 3, illustrating the effects of the initial subsurface states, SL and  $T_e$ , on prediction of the 2006 El Niño event. Different from the discrepancy induced by the initial surface conditions, the Assim\_SST + SL hindcast had an obviously stronger variation in SL than that of Assim\_SST, which is associated with the propagation of two downwelling Kelvin waves during the forecast process. As shown in Fig. 3c and b, the early warming at both the surface and subsurface layers over the eastern equatorial Pacific in the Assim\_SST + SL hindcast corresponds to a weak Kelvin downwelling wave initiated east of the dateline before 2005 (Fig. 3a). An additional stronger Kelvin downwelling wave propagated eastbound in approximately January 2006 in the equatorial western Pacific (Fig. 3a and b) and reached the eastern tropical Pacific in mid-2006. These two eastward downwelling propagations resemble the observed events and agree with the timing of the large deepening anomalies of the thermocline depth in the equatorial eastern Pacific (not shown), the details of which have been reported by Hackert et al. (2007). Finally, a 0.8 °C warmer El Niño condition at the end of 2006 was predicted when considering the accuracy of the initial subsurface condition, and much stronger coupled air–sea interaction and vertical mixing induced by the SL assimilation were evident during the second half of 2006 when compared to those induced by the SST assimilation.

As a result, the inclusion of more reasonable and accurate initial conditions during the 2006–2007 El Niño forecast process can lead to better predictions. Figure 4 compares the hindcasts initiated from the three initialization schemes for the 2006–2007 El Niño episode. The hindcasts initialized by the Assim\_SST + SL scheme could successfully predict the onset, development, and decay of 2006 El Niño at all times prior to the event, although some small errors remained in the onset and magnitude of the 2006–2007 El Niño event when forecasted nine months ahead. The hindcasts

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

subsurface field than by that of the surface field (Zheng et al., 2007; Zheng and Zhu, 2008), suggesting that advanced data assimilation methodology combined with accurate subsurface information should be used in conjunction with coupled forecasts to extend the lead time and accuracy of ENSO forecast systems (e.g., Zhu et al., 2012).

5 Analysis of the results show that the initial conditions having optimal subsurface structures such as SL and  $T_e$  fields from the EnKF analysis mainly contributed to improving the intensity prediction in the growth of 2006–2007 El Niño, although an accurate initial surface state such as SST or wind stress fields can help to easily trigger a warming tendency for predicting the 2006–2007 El Niño event. It should be noted that this work is  
10 limited to only one event, and it is difficult to make generic statements about all El Niño events. However, the conceptually simple and robust, technique used here will help coupled modelers to isolate the role of the initial state of the ocean and to evaluate the conditions favorable for improving El Niño predictions.

15 *Acknowledgements.* This work was supported by the National Basic Research Program of China (Grant Nos. 2012CB417404 and 2012CB955202), WPOS (Grant No. XDA10010405), and the Natural Science Foundation of China (Grant No. 41176014).

## References

- Alves, O., Balmaseda, M., Anderson, D., and Stockdale, T.: Sensitivity of dynamical seasonal forecasts to ocean initial conditions, *Q. J. Roy. Meteor. Soc.*, 130, 647–668, 2004.
- 20 Ashok, K. and Yamagata, T.: Climate change: the El Niño with a difference, *Nature*, 461, 481–484, 2009.
- Barnston, A. G., Tippett, M. K., L'Heureux, M. L., Li, S., and DeWitt, D. G.: Skill of real-time seasonal ENSO model predictions during 2002–2011: is our capability increasing?, *B. Am. Meteorol. Soc.*, 93, 631–651, 2012.
- 25 Evensen, G.: Sampling strategies and square root analysis schemes for the EnKF, *Ocean Dynam.*, 54, 539–560, 2004.
- Evensen, G.: *Data Assimilation: the Ensemble Kalman Filter*, 2nd edn., Springer, Berlin, 2009.

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Hackert, E., Ballabrera-Poy, J., Busalacchi, A. J., Zhang, R.-H., and Murtugudde, R.: Role of the initial ocean state for the 2006 El Niño, *Geophys. Res. Lett.*, 34, L09605, doi:10.1029/2007GL029452, 2007.

Ji, M. and Leetmaa, A.: Impact of data assimilation on ocean initialization and El Niño prediction, *Mon. Weather Rev.*, 125, 742–753, 1997.

Ji, M., Reynolds, R. W., and Behringer, D. W.: Use of TOPEX/Poseidon sea level data for ocean analyses and ENSO prediction: some early results, *J. Climate*, 13, 216–231, 2000.

Jin, E. K., Kinter III, J. L., Wang, B., Park, C.-K., Kang, I.-S., Kirtman, B. P., Kug, J.-S., Kumar, A., Luo, J.-J., Schemm, J., Shukla, J., and Yamagata, T.: Current status of ENSO prediction skill in coupled ocean–atmosphere models, *Clim. Dynam.*, 31, 647–664, 2008.

Keenlyside, N. and Kleeman, R.: Annual cycle of equatorial zonal currents in the Pacific, *J. Geophys. Res.*, 107, doi:10.1029/2000JC000711, 2002.

Keenlyside, N., Latif, M., Botzet, M., Jungclaus, J., and Schulzweida, U.: A coupled method for initialising ENSO forecasts using SST, *Tellus A*, 57, 340–356, 2005.

Kirtman, B. P., Shukla, J., Balmaseda, M., Graham, N., Penland, C., Xue, Y., and Zebiak, S. E.: Current status of ENSO forecast skill, A report to the Climate Variability and Predictability (CLIVAR) Numerical Experimentation Group (NEG), CLIVAR Working Group on Seasonal to Interannual Prediction, 2002.

Latif, M., Anderson, D., Barnett, T., Cane, M., Kleeman, R., Leetmaa, A., O'Brien, J., Rosati, A., and Schneider, E.: A review of the predictability and prediction of ENSO, *J. Geophys. Res.*, 103, 14375–14393, 1998.

McCreary, J. P.: A linear stratified ocean model of the equatorial undercurrent, *Philos. T. R. Soc. Lond.*, 298, 603–635, 1981.

McPhaden, M. J.: Evolution of the 2006–2007 El Niño: the role of intraseasonal to interannual time scale dynamics, *Adv. Geosci.*, 14, 219–230, doi:10.5194/adgeo-14-219-2008, 2008.

McPhaden, M. J., Zebiak, S. E., and Glantz, M. H.: ENSO as an integrating concept in earth science, *Science*, 314, 1739–1745, 2006.

Neelin, J. D., Battisti, D. S., Hirst, A. C., Jin, F.-F., Wakata, Y., Yamagata, T., and Zebiak, S. E.: ENSO theory, *J. Geophys. Res.*, 103, 14262–14290, 1998.

Palmer, T. N.: A nonlinear dynamical perspective on model error: a proposal for non-local stochastic-dynamic parametrization in weather and climate prediction models, *Q. J. Roy. Meteor. Soc.*, 127, 279–304, 2001.

## Roles of initial ocean surface and sub-surface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Picaut, J., Hackert, E., Busalacchi, A. J., Murtugudde, R., and Lagerloef, G. S. E.: Mechanisms of the 1997–1998 El Niño–La Niña, as inferred from space-based observations, *J. Geophys. Res.*, 107, 3037, doi:10.1029/2001JC000850, 2002.

Sakov, P. and Oke, P. R.: Implications of the form of the ensemble transformations in the ensemble square root filters, *Mon. Weather Rev.*, 136, 1042–1053, 2008.

Smith, T. M., Reynolds, R. W., Peterson, T. C., and Lawrimore, J.: Improvements to NOAA's historical merged land–ocean surface temperature analysis (1880–2006), *J. Climate*, 21, 2283–2296, 2008.

Yang, S.-C., Rienecker, M., and Keppenne, C.: The impact of ocean data assimilation on seasonal-to-interannual forecasts: a case study of the 2006 El Niño event, *J. Climate*, 23, 4080–4095, 2010.

Zebiak, S. E. and Cane, M. A.: A model El Niño–Southern Oscillation, *Mon. Weather Rev.*, 115, 2262–2278, 1987.

Zhang, R.-H., Zebiak, S. E., Kleeman, R., and Keenlyside, N.: Retrospective El Niño forecast using an improved intermediate coupled model, *Mon. Weather Rev.*, 133, 2777–2802, 2005.

Zheng, F. and Zhu, J.: Balanced multivariate model errors of an intermediate coupled model for ensemble Kalman filter data assimilation, *J. Geophys. Res.*, 113, C07002, doi:10.1029/2007JC004621, 2008.

Zheng, F. and Zhu, J.: Coupled assimilation for an intermediated coupled ENSO prediction model, *Ocean Dynam.*, 60, 1061–1073, doi:10.1007/s10236-010-0307-1, 2010.

Zheng, F., Zhu, J., Zhang, R.-H., and Zhou, G.-Q.: Ensemble hindcasts of SST anomalies in the tropical Pacific using an intermediate coupled model, *Geophys. Res. Lett.*, 33, L19604, doi:10.1029/2006GL026994, 2006.

Zheng, F., Zhu, J., and Zhang, R.-H.: The impact of altimetry data on ENSO ensemble initializations and predictions, *Geophys. Res. Lett.*, 34, L13611, doi:10.1029/2007GL030451, 2007.

Zhu, J., Huang, B., Marx, L., Kinter III, J. L., Balmaseda, M. A., Zhang, R.-H., and Hu, Z.-Z.: Ensemble ENSO hindcasts initialized from multiple ocean analyses, *Geophys. Res. Lett.*, 39, L09602, doi:10.1029/2012GL051503, 2012.

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu

**Table 1.** Summary of the initialization scheme design.

Initialization Scheme	Assimilated Data		Description
	SST	SL	
Assim_SST	✓		Optimizing the initial surface conditions (i.e., SST and wind stress)
Assim_SL		✓	Updating the initial subsurface states (i.e., SL and $T_e$ )
Assim_SST + SL	✓	✓	Improving the accuracy of both the initial surface and subsurface states

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Roles of initial ocean surface and sub-surface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu




**Figure 1.** Initial conditions of anomalous sea surface temperature (SST; top row), sea level (SL; middle row), and forecasted SST anomalies (bottom row) from the (a) observations, (b) Assim\_SST results, (c) Assim\_SL results, and (d) Assim\_SST + SL results, respectively. The initial fields are from December 2005, and the 12 month forecasts start in January 2006.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Roles of initial ocean surface and sub-surface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu



**Figure 2.** Initial and forecasted differences between the Assim\_SST + SL hindcast and the Assim\_SL hindcast for **(a)** sea surface temperature (SST), **(b)** zonal wind stress (Taux), **(c)** sea level (SL), and **(d)** temperature of the subsurface water entrained into the mixed-layer ( $T_e$ ), illustrating the effects of the initial surface states, SST and Taux on predicting the 2006–2007 El Niño event. The dashed red line in **(c)** represents the propagation of the downwelling Kelvin wave. The contour interval is  $0.2^\circ\text{C}$  in **(a)**,  $0.08\text{ dyn cm}^{-2}$  in **(b)**,  $0.8\text{ cm}$  in **(c)**, and  $0.1^\circ\text{C}$  in **(d)**.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Roles of initial ocean surface and subsurface states on successfully predicting 06/07 El Niño


F. Zheng and J. Zhu



**Figure 3.** Initial and forecasted differences between the *Assim\_SST + SL* hindcast and *Assim\_SST* hindcast for **(a)** sea level (SL), **(b)** temperature of the subsurface water entrained into the mixed-layer ( $T_e$ ), **(c)** sea surface temperature (SST), and **(d)** zonal wind stress ( $Taux$ ), illustrating the effects of the initial subsurface states, SL and  $T_e$  on predicting the 2006–2007 El Niño event. The dashed red line in **(a)** represents the propagation of the downwelling Kelvin wave. The contour interval is 1.0 cm in **(a)**, 0.2 °C in **(b)** and **(c)**, and 0.04 dyn cm<sup>-2</sup> in **(d)**.

## Roles of initial ocean surface and sub-surface states on successfully predicting 06/07 El Niño

F. Zheng and J. Zhu



**Figure 4.** Comparisons of the hindcast results for the 2006–2007 El Niño event. The thick black curves are the observed Niño 3.4 sea surface temperature (SST) anomalies, and the thin curves of red, green, and blue are the predictions initialized by the Assim\_SST + SL, Assim\_SST, and Assim\_SL assimilation results, respectively. **(a)**, **(b)**, and **(c)** are hindcasts that started from 12, 9, and 6 months, respectively, ahead of the peak of the 2006–2007 El Niño event.