

Abstract

We document dense water formation (DWF) throughout the Adriatic shelf and coastal area in January/February 2012, resulting in record-breaking densities observed during and after the event. The unprecedented dense water generation was preconditioned by a dry and warm year which resulted in a significant reduction of coastal freshwaters, superimposed on a long-term basin-wide salinity increase. The final event that triggered the DWF was an extended period of cold weather with strong and severe winds. Record-breaking potential density anomalies (above 30 kg m^{-3}) were measured at several DWF sites. Accumulated surface net heat and water losses in some coastal regions exceeded 1.5 GJ m^{-2} and 250 kg m^{-2} over 21 days, respectively. Excessiveness, importance of shelf-type DWF, effects on the thermohaline circulation and deep aquatic systems, and connection with climate change are discussed.

1 Introduction

Aside from the open-ocean convection type of dense water formation (DWF), which has been early recognized to be substantial in driving thermohaline circulation in global ocean and deep basins, the role of DWF over a shelf in changes of deep basins has not been perceived as important until recently (Ivanov et al., 2004; Allen and Durrieu de Madron, 2009; Pattiaratchi et al., 2011). This applies particularly to mid-latitude large and deep basins, such as the Mediterranean Sea, where deep thermohaline circulation is driven by extensive wintertime heat losses and wind mixing at several DWF sites (Robinson et al., 2003). Although extensively investigated at the Mediterranean sub-basin and process scales (Pasqual et al., 2010), the contribution of the shelf-generated dense water has generally been underrated in the Mediterranean-scale studies, especially within long-term and climate studies (Somot et al., 2006). This is partly a result of still too low resolution of coastal bathymetry and strong temporal and spatial variability of the atmospheric forcing in shelf regions (Béranger et al., 2010).

OSD

9, 3701–3721, 2012

Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

et al., 2010). Conversely, the changing properties of outflowing AdDW eventually modify the upper-layer circulation in the Ionian and the pathways of water masses, thus impacting the entire Levantine basin (Gačić et al., 2011).

Multi-decadal monitoring efforts along the DWF area and several campaigns that captured very dense bottom currents (Supić and Vilibić, 2006) documented maximum potential density anomaly (PDA – σ_θ) in the Adriatic not exceeding 30 kg m^{-3} . However, it seems that higher densities may be reached quite close to the coast, during severe winters occurring over centurial timescales. Namely, Vatova (1929) documented the temperature of 3.95°C and salinity equaling 38.15 at 22 m depth off Koper (Gulf of Trieste), during severe winter of 1929, resulting in PDA of 30.30 kg m^{-3} . Harsh winter conditions caused mass mortality of benthic organisms at the time. Still, this almost 100 yr old measurement is the only historical record with such a high density in the North Adriatic. The record was broken during the long and severe bora episode in the winter of 2012, preconditioned by a very dry year. During this recent episode the shelf and coastal DWF occurred not only at the classical DWF site but also in a number of Eastern Adriatic coastal channels and bays, where DWF has not been documented to this day due to the large freshwater load. This paper is the first documentation of the DWF that occurred in the North and Middle Adriatic during the winter of 2012. The study includes an assessment of the preconditioning and atmospheric forcing reproduced by operational atmospheric and ocean models, and it raises a discussion on different aspects of the observed extreme event.

2 Data and methods

2.1 Field campaigns and data processing

A number of oceanographic field campaigns were carried out in the coastal and open Adriatic waters between February and April 2012, taking CTD profiles by different probes (Seabird SBE 19 and SBE 25, Idronaut 316, Sea&Sun MSS90), all of them

Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dynamics and a full set of physics parameterizations (Tudor and Ivatek-Šahdan, 2010). It is initialized daily from the 6 h ALADIN forecast run in 8 km resolution using the scale selective digital filter initialization (Termonia, 2008). SST in the model provided by the global atmospheric model ARPEGE and does not change during the 24 h model run.

5 Turbulent fluxes are computed as turbulent exchange between the lowest model level (around 17 m above the sea surface) and the sea surface. The turbulent exchange coefficients are resolved using prognostic turbulent kinetic energy (TKE) values.

The second COSMO model (www.cosmo-model.org) is a limited-area nonhydrostatic atmospheric prediction model with 7 km horizontal resolution and 72-h forecast range, being operational at Hydro-Meteo-Clima Service (SIMC) of the Emilia Romagna Regional Agency for Protection and Environment (ARPA-EMR). The Regional Ocean Modeling System (ROMS, <http://www.myroms.org>) is a primitive equation, finite differences, hydrostatic, free surface model, tides included. The utilized Adriatic application (described in Russo et al. (2009) and operational at DISVA in collaboration with SIMC and CNR-ISMAR) has a constant resolution of 2 km across the entire domain and 20 stretched vertical sigma levels. Forty-nine rivers and karstic springs (in the eastern coastal area) were also included as sources of mass and momentum using available daily discharges for the Po River and monthly climatological values for other sources. Radiation condition is imposed at the Strait of Otranto with a nudging of mean daily forecast of temperature, salinity, baroclinic currents from the general circulation model of the Mediterranean Ocean Forecasting System (Oddo et al., 2009). Fluxes through the air-sea interface were calculated using the COARE 3.0 bulk flux algorithms (Fairall et al., 2003), with sea surface temperature (SST) from ROMS and short wave radiation, wind, air temperature, humidity, cloud cover and atmospheric pressure from the COSMO-I7 atmospheric model.

Exceptional dense water formation on the Adriatic shelf in the winter of 2012H. Mihanović et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3 Results and discussion

Maximum PDA records and tabulated respective PTs, salinities and dates of measurements taken in the North and Middle Adriatic between February and April 2012 (during and after the extreme cold event and severe bora outbreak) are shown in Fig. 1. The bora episode lasted for about 3 weeks in coastal Eastern Adriatic region, between 25 January and 14 February 2012. Wind was strong to severe from ENE–NE, blowing occasionally with hurricane force. The maximum PDAs were measured throughout the Gulf of Trieste, with absolute maximum at TS station sampled on 15 February (30.59 kg m⁻³ at 18 m depth; the respective PT and salinity were 4.24 °C and 38.56). During the field study carried out in the Gulf between 14 and 16 February 2012, PDA surpassed 30.50 kg m⁻³ at 25 CTD stations out of 44. The absolute temperature minimum below 20 m was observed at 24 m depth on the Paloma pillar, in the central part of the Gulf (3.9 °C), in the mornings of 13 and 15 February 2012. This temperature record is quite similar to the bottom temperature observed by Vatova (1929) in the winter of 1929 (3.95 °C at 22 m).

Slightly lower PDAs were measured at the AA oceanographic platform on 11 February (30.35 kg m⁻³). The platform is located at the margin of the bora-driven Northern Adriatic gyre, in which a peak of DWF is normally located (Boldrin et al., 2009). Such densities are much higher than those previously recorded in the area over an extended period of time (Supić and Vilibić, 2006), driven both by extremely high salinities (> 38.5) and very low temperatures (< 6.5 °C). Another PDA maximum was measured at 100-m deep Pag Channel station (PC), located deep inside Croatian inner waters. Exceptionally high salinities (around 38.77) were recorded at the station and these salinities are even higher than the average value documented for the South Adriatic (Artegiani et al., 1997). Finally, extremely high salinities, low PTs and associated high PDAs were recorded along the entire Eastern Adriatic coastal area northwest from Split, all above 29.55 kg m⁻³. Such values have never been documented before in this part of the Adriatic (e.g. Zore-Armanda et al., 1991).

Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Spatial distribution of modelled fluxes, represented in Fig. 5 as a cumulative energy and mass loss during the bora episode (25 January–14 February 2012), quantifies the amount of energy and mass taken from the sea and indicates jets with extreme energy losses due to cold and severe bora wind. The energy loss (Fig. 5a) exceeded 1.5 GJm⁻² over the maximum bora jet along the eastern shore (off Senj), which is equivalent to a cooling rate of about 4.5 °C over the whole event, if box-model heat balance approach over 80 m deep ocean (average sea depth over the bora jets areas in the inner Croatian waters) is used (Gill, 1982). Naturally, similar energy losses over smaller and shallower embayments, e.g. Pag Bay (40 m average depth), would multiply the cooling of the ocean. Lower cooling rates over shallower Northern Adriatic would yield a similar temperature decrease. The energy loss of 0.7 GJm⁻² modelled over 30–40 m deep TL embayment would decrease the temperature for about 5 °C. Similar rates were estimated for the Gulf of Trieste and confirmed by temperature measurements from the buoys in the Gulf. Lower heat losses occurred over the DWF open-convection area in the South Adriatic, where bora did not blow during the whole period and therefore relatively modest cooling and vertical mixing were observed (up to 500 m).

In addition to the cooling, cumulative E-P water loss (Fig. 5b) during the event surpassed 250 kgm⁻² over the maximum bora jet, being equivalent to a salinity increase of 0.12 over the 80 m deep inner Croatian waters. Cumulative water losses were 2–3 times lower over classical DWF site in the Northern Adriatic, resulting in a slightly lower salinity increase rate (about 0.1). By using a box model, PDA increase can be estimated to around 0.95 kgm⁻³ in the inner Croatian waters.

The volume of the generated NAdDW exceeding the density threshold of 29.5 kgm⁻³ (Artegiani et al., 1997) equalled 4250 km³, estimated from the COSMO/ROMS numerical model. The NAdDW encompassed the most of the northern and large part of the Middle Adriatic. By assuming that the associated dense water current (flowing south-east) lasts for 3 months, one may reach a preliminary estimate of average transport of 0.55 Sv, which is an order of magnitude larger than during normal DWF years (Vilibić and Supić, 2005).

on 26 April 1956, where PT and salinity at 240 m equalled to 9.27°C and 38.31, respectively, resulting in a PDA of 29.67 kg m⁻³. One should note that both past extreme DWFs (in 1929 and 1956) were characterized by much lower salinities, as preconditioning was presumably different, with much higher precipitation rates and river runoffs and cooler climate preceding the DWF.

The observed difference in temperature and salinity over centurial timescales may be associated with ongoing climate changes, observable in a positive trend of the Adriatic salinity and upper-layer temperature (Vilibić et al., 2012), driven by positive E-P-R and heat flux trends observed over the Adriatic. Climate projection for the end of 21st century gives additional warming of surface Adriatic waters by about 2.5° C and salinity increase of about 0.5 (Tsimplis et al., 2008), changing the preconditioning of the DWF. This will presumably result in more saline and warmer dense waters, replacing present deep Adriatic and Mediterranean waters. However, the changes in coastal regions are found and projected to be more rapid: the decrease in freshwater coastal input in the Adriatic, resulting in salinity and density trends higher in coastal regions than in the open Adriatic (Vilibić et al., 2012), may weaken Adriatic-Ionian thermohaline circulation and therefore change the BiOS variability.

On the other hand, atmospheric blocking situations and intensity of cold air outbreaks over Southern Europe are projected to increase in the 21st century (Vavrus et al., 2006). This may result in more frequent occurrence of prolonged DWF episodes like in 2012, which may strengthen BiOS and the Adriatic-Ionian thermohaline cell. It seems that interplay between prolonged DWF during winter blocking situation and “normal” winters in the future climate will determine the strength of the thermohaline circulation, although regional climate models project that the circulation could become shallower (Somot et al., 2006). These changes are quite important, as the DWF brings oxygen to deep ocean layers, necessary especially within biodiversity niches such is Jabuka Pit, but also important for the whole deep Mediterranean. Finally, they will influence internal processes in the Ionian Sea and the BiOS, thus affecting circulation patterns

Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and pathways of water masses, and possibly impacting the future interplay in DWF between the Adriatic and Aegean Seas.

Acknowledgements. We are indebted to all the oceanographers, crew members and technicians who collected and processed the observations. The data collected from the ARGO floats were made freely available by the International Argo Program and the national programs that contribute to it. We are grateful to A. Coluccelli (DISVA, Università Politecnica delle Marche, Ancona, Italy) for ROMS operational setup and maintenance and to M. Deserti and A. Valentini (SIMC-ARPA-EMR, Bologna, Italy) for the COSMO forecasts. The support for the study was received through the Ministry of Science, Education and Sports of the Republic of Croatia (grants 001-0013077-1122, 119-1193086-3085, 119-1191189-1228 and 004-1193086-3036). A. Bergamasco and A. Boldrin acknowledge the partial support received by the Veneto Region (Regional Law no. 15, 12 July 2007, Projects SISOE – Integrated System for Oceanographic and Ecological monitoring of the Adriatic Sea and MARINA), while S. Carniel is grateful to the FIRB Project support. V. Malačič is indebted to the group performing the national monitoring program (under the EU Water Framework Directive), funded by the Slovenian Agency of Environment, and to the research program funded by the Ministry of Higher Education and Technology of Slovenia (contract no. P1-0237). M. Celio is indebted to the marine coastal waters monitoring program of the Friuli Venezia Giulia Region under the EU Water Framework Directive 2000/60/CE and D.M. 260/10.

References

- Allen, S. E. and Durrieu de Madron, X.: A review of the role of submarine canyons in deep-ocean exchange with the shelf, *Ocean Sci.*, 5, 607–620, doi:10.5194/os-5-607-2009, 2009.
- Artegiani, A., Bregant, D., Paschini, E., Pinardi, N., Raicich, F., and Russo, A.: The Adriatic Sea general circulation. Part I: Air-sea interactions and water mass structure, *J. Phys. Oceanogr.*, 27, 1492–1514, 1997.
- Béranger, K., Drillet, Y., Housais, M. N., Testor, P., Bourdallé-Badie, R., Alhammoud, B., Bozec, A., Mortier, L., Bouruet-Aubertot, P., and Crépon, M.: Impact of the spatial distribution of the atmospheric forcing on water mass formation in the Mediterranean Sea, *J. Geophys. Res.*, 115, C12041, doi:10.1029/2009JC005648, 2010.

Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Bergamasco, A., Oguz, T., and Malanotte-Rizzoli, P.: Modeling dense water mass formation and winter circulation in the Northern and Central Adriatic Sea, *J. Marine Syst.*, 20, 279–300, 1999.
- Boldrin, A., Carniel, S., Giani, M., Marini, M., Bernardi Aubry, F., Campanelli, A., Grilli, F., and Russo, A.: Effects of bora wind on physical and biogeochemical properties of stratified waters in the Northern Adriatic, *J. Geophys. Res.*, 114, C08S92, doi:10.1029/2008JC004837, 2009.
- Civitaresse, G., Gačić, M., Lipizer, M., and Eusebi Borzelli, G. L.: On the impact of the Bi-modal Oscillating System (BIOS) on the biogeochemistry and biology of the Adriatic and Ionian Seas (Eastern Mediterranean), *Biogeosciences*, 7, 3987–3997, doi:10.5194/bg-7-3987-2010, 2010.
- Edson, J. B.: Review of air-sea transfer processes, in: Proceedings on ECMWF Workshop on Ocean-Atmosphere Interactions, ECMWF, Reading, 7–24, 10–12 November 2008, 2009.
- Fairall, C. W., Bradley, E. F., Hare, J. E., Grachev, A. A., and Edson, J. B.: Bulk parameterization of air-sea fluxes: updates and verification for the COARE algorithm, *J. Climate*, 16, 571–591, 2003.
- Gačić, M., Civitaresse, G., Miserocchi, S., Cardin, V., Crise, A., and Mauri, E.: The open-ocean convection in the Southern Adriatic: a controlling mechanism of the spring phytoplankton bloom, *Cont. Shelf Res.*, 22, 1897–1908, 2002.
- Gačić, M., Eusebi Borzelli, G. L., Civitaresse, G., Cardin, V., and Yari, S.: Can internal processes sustain reversals of the ocean upper circulation?: the Ionian Sea example, *Geophys. Res. Lett.*, 37, L09608, doi:10.1029/2010GL043216, 2010.
- Gačić, M., Civitaresse, G., Eusebi Borzelli, G. L., Kovačević, V., Poulain, P.-M., Theocharis, A., Menna, M., Catucci, A., and Zarokanellos, N.: On the relationship between the decadal oscillations of the Northern Ionian Sea and the salinity distributions in the Eastern Mediterranean, *J. Geophys. Res.*, 116, C12002, doi:10.1029/2011JC007280, 2011.
- Gill, A. E.: *Atmosphere-Ocean Dynamics*, Academic Press, Orlando, 662 pp., 1982.
- Grisogono, B. and Belušić, D.: A review of recent advances in understanding the meso- and microscale properties of the severe Bora wind, *Tellus A*, 61, 1–16, 2009.
- Ivanov, V. V., Shapiro, G. I., Huthnance, J. M., Aleynik, D. L. and Golovin, P. N.: Cascades of dense water around the world ocean, *Prog. Oceanogr.*, 60, 47–98, 2004.
- Jeffries, M. A. and Craig, M. L.: A climatology of the Northern Adriatic Sea's response to bora and river forcing, *J. Geophys. Res.*, 112, C03S02, doi:10.1029/2006JC003664, 2007.

Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Somot, S., Sevault, F., and Déqué, M.: Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model, *Clim. Dynam.*, 27, 851–879, 2006.

Supić, N. and Vilibić, I.: Dense water characteristics in the Northern Adriatic in the 1967–2000 interval with respect to surface fluxes and Po River discharges, *Estuar. Coast. Shelf S.*, 66, 580–593, 2006.

Termonia, P.: Scale selective digital filter initialization, *Mon. Weather Rev.*, 136, 5246–5255, 2008.

Tsimplis, M. N., Marcos, M., and Somot, S.: 21st century Mediterranean sea level rise: steric and atmospheric pressure contributions from a regional model, *Global Planet. Change*, 63, 105–111, 2008.

Tudor, M. and Ivatek-Šahdan, S.: The case study of bura of 1st and 3rd February 2007, *Meteorol. Z.*, 19, 2867–2882, 2010.

Vatova, A.: Sui minimi termici verificatisi nell'alto Adriatico nel febbraio e nel marzo 1929 e loro effetti sull'ittiofauna, *Regio Comitato Talassografico Italiano, Memoria*, 157, 1–8, 1929 (in Italian).

Vavrus, S., Walsh, J. E., Chapman, W. L., and Portis, D.: The behavior of extreme cold air outbreaks under greenhouse warming, *Int. J. Climatol.*, 26, 1133–1147, 2006.

Vilibić, I. and Supić, N.: Dense water generation on a shelf: the case of the Adriatic Sea, *Ocean Dynam.*, 55, 403–415, 2005.

Vilibić, I., Šepić, J., and Proust, N.: Observational evidence of a weakening of thermohaline circulation in the Adriatic Sea, *Clim. Res.*, doi:10.3354/cr01128, in press, 2012.

Zore-Armanda, M.: Les masses d'eau de la mer Adriatique, *Acta Adriat.*, 10, 5–88, 1963.

Zore-Armanda, M., Bone, M., Dadić, V., Morović, M., Ratković, D., Stojanoski L., and Vukadin, I.: Hydrographic properties of the Adriatic Sea from 1971 through 1983, *Acta Adriat.*, 32, 1–547, 1991.




Fig. 1. Adriatic Sea bathymetry, with selected stations at which CTD samplings were carried out between February and April 2012 (main panel – the whole Adriatic; smaller inset – zoom of the Eastern Adriatic coastal area). Maximum values of potential density anomalies (PDA, in kg m^{-3}) observed at these locations are indicated adjacent to station names. Respective sampling dates, depths, salinities and potential temperatures (PT) are shown in an inserted table. Continuous measurements of meteorological and oceanographic parameters were available at Acqua Alta platform (AA). The green line in smaller inset indicates CTD transect used in Fig. 2c. The main path of the North Adriatic Dense Water (NAdDW) appears within blue dashed lines in the main panel. Presumed pathways of dense water generated along Eastern Adriatic coast are denoted by blue arrows in smaller inset.

Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.




Fig. 2. Vertical profiles of potential temperature, practical salinity and potential density anomaly at selected stations along the Eastern Adriatic coast: **(a)** JP on 17 September 2011, 23 February 2012 and 27 April 2012, **(b)** BL on 28 February 2012, TL on 29 February 2012 and PC on 29 March 2012.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Distribution of **(a)** potential temperature (PT, in $^{\circ}\text{C}$), **(b)** practical salinity and **(c)** potential density anomaly (PDA, in kg m^{-3}) measured in the Gulf of Trieste between 14 and 16 February 2012. The mouth of the Isonzo River is at the northwestern part of the transect. See the smaller inset in Fig. 1 for transect position.




Fig. 4. (a) Time series of turbulent heat fluxes estimated for AA, together with modelled ones computed by ALADIN/HR and COSMO/ROMS models at the grid point nearest to AA, for the period between 25 January and 14 February 2012, and time series of surface (2 m) and bottom (13 m) (b) potential temperature (PT), (c) practical salinity and (d) potential density anomaly (PDA) during the same period.

Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Exceptional dense water formation on the Adriatic shelf in the winter of 2012

H. Mihanović et al.




Fig. 5. Cumulative **(a)** heat and **(b)** water gain/loss between 25 January and 14 February 2012 as estimated by ALADIN/HR model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion