



This discussion paper is/has been under review for the journal Ocean Science (OS).
Please refer to the corresponding final paper in OS if available.

Arctic Ocean circulation and variability – advection and external forcing encounter constraints and local processes

B. Rudels^{1,*}

¹Department of Physics, University of Helsinki and Finnish Meteorological Institute, Helsinki, Finland

*Invited contribution by B. Rudels, recipient of the Fridtjof Nansen Medal 2011.

Received: 27 October 2011 – Accepted: 14 November 2011 – Published: 1 December 2011

Correspondence to: B. Rudels (bert.rudels@fmi.fi)

Published by Copernicus Publications on behalf of the European Geosciences Union.

OSD

8, 2313–2376, 2011

Arctic Ocean
Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

Discussion Paper | Discussion Paper | Discussion Paper

OSD

8, 2313–2376, 2011

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



The first hydrographic data from the Arctic Ocean, the section from the Laptev Sea to the passage between Greenland and Svalbard obtained by Nansen on the drift by Fram 1893–1896, aptly illustrate the main features of Arctic Ocean oceanography and indicate possible processes active in transforming the water masses in the Arctic Ocean. Many, perhaps most, of these processes were identified already by Nansen, who put his mark on almost all subsequent research in the Arctic Ocean. Here we shall revisit some key questions and follow how our understanding has evolved from the early 20th century to present. What questions, if any, can now be regarded as solved and which remain still open? Five different but connected topics will be discussed: (1) The low salinity surface layer and the storage and export of freshwater. (2) The vertical heat transfer from the Atlantic water to sea ice and to the atmosphere. (3) The circulation and mixing of the two Atlantic inflow branches. (4) The formation and circulation of deep and bottom waters in the Arctic Ocean. (5) The exchanges through Fram Strait. Foci will be on the potential effects of increased freshwater input and reduced sea ice export on the freshwater storage and residence time in the Arctic Ocean, on the deep waters of the Makarov Basin and on the circulation and relative importance of the two inflows, over the Barents Sea and through Fram Strait, for the distribution of heat in the intermediate layers of the Arctic Ocean.

20 1 Introduction

25

In the 19th century the high Arctic was an unknown part of the world. Was it an ocean or were still large landmasses to be found? Was it covered with ice or was open water present beyond the forbidding ice fields and hummocks that in the 16th and 17th centuries had prevented all expeditions to navigate the Northwest and the Northeast Passages? Frobisher, Davis and Baffin explored Davis Strait and Baffin Bay and Willem Barents the Barents Sea and the waters west of Svalbard in search of the Northwest

and Northeast Passages. Hudson tried to find both. All efforts were in vain and eventually these navigational goals were replaced by hunting of whales and walruses. In the 17th century a Dutch whaling industry was established on Svalbard and the northern North America was explored and exploited by British and French companies like the 5 Hudson Bay Company.

In the 18th century Russian explorers charted the Siberian coast and Vitus Bering explored the passage between North America and Asia, Bering Strait, but it was not until the beginning of the 19th century that renewed efforts were made to discover a navigational route north of North America. The expeditions of J. Ross (1818, 1829–1833) and

10 W. E. Parry (1819–1820, 1821–1822, 1824–1825) did not succeed. J. Phipps (1773), W. Scoresby (1806), D. Buchan and J. Franklin (1816) and also Parry (1827) tried to sail north between Greenland and Svalbard but were stopped by the ice. Parry used sledges to go farther north and reached ($82^{\circ}45'$) but had to return. The ice was drifting southward with a speed comparable to the progress over the ice and the northward 15 advance was close to zero. The disappearance of the large expedition led by Franklin (1845) and the subsequent search for its fate added much information about the straits and islands in the now Canadian Arctic Archipelago but did not lead to the discovery of a navigable sea route connecting the North Atlantic to the North Pacific north of America. Ice was blocking the way.

20 In spite of these setbacks a vision of an ice-free ocean beyond the pack ice arose in the late 18th and early part of the 19th century. M. Lomonosov, L. A. de Bougainville and S. Engel were among the first proponents of an ice-free polar ocean. Engel (1772) thought that since sea ice contains little salt only freshwater could freeze and once a ship passed through the girdle of low salinity water created by the river runoff a navigable 25 ocean would be found. D. Barrington, vice-president of the Royal Society, believed in an open Polar Ocean and was promoting Phipps' expedition in 1773. The idea was getting stronger with time and in the mid 19th century it found its principal spokesman in the German geographer Augustus Petermann. Petermann's arguments for an ice free polar ocean was partly that open water has been observed north of

Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Siberia also in winter beyond the frozen coastal areas, but also that the ocean currents, especially the Gulf Stream, were transporting warm water northward to the Arctic. This heat would be sufficient to prevent ice to form and keep an ocean close to the North Pole free of ice (Petermann, 1865) (Fig. 1).

- 5 The most favourable routes to penetrate northward into this area of open water would be to follow the warm currents, the Norwegian Atlantic Current and the West Spitsbergen Current between Greenland and Svalbard. A second possibility would be east of Svalbard, where a part of the Norwegian Atlantic Current flows into the Barents Sea and then continues north. Petermann was involved in the planning of the German North
- 10 Pole expeditions led by C. Koldeway (1868, 1869–1870) and was also giving advice to the Austrian Tegetthoff expedition (1872–1874) led by K. Weyprecht and J. Payer. Both these expeditions ended by ships getting caught in the ice and abandoned. During the same period Great Britain and the United States were, after the search for Franklin, who had disappeared in an attempt to navigate the Northwest Passage, trying to reach
- 15 the open Polar Ocean west of Greenland. The expeditions by E. K. Kane (1853–1855), I. Hayes (1860–1861), C. Hall (1871–1873) and G. Nares (1875–1876) penetrated north between Greenland and Ellesmere Island, the Nares Strait, into the Arctic Ocean to finally be stopped by the ice.

The unknown area, where an ice-free polar ocean could exist, diminished gradually

20 and in 1879 the vessel Jeanette commanded by G. Delong sailed north to Bering Strait to look for A.E. Nordenskiöld and Wega. However, as Delong reached Bering Strait, Wega had already completed the Northeast Passage and passed south into the North Pacific. Delong continued northward through Bering Strait to try to reach the North Pole from the Pacific sector. Jeanette got caught in the ice the first winter and then

25 again the second winter and the ship had to be abandoned in 1881. A few members of the crew made it back but Delong was not among those.

Three years after the wreckage of Jeanette parts from the ship were found at Julianehåb on the southwest coast of Greenland. This discovery indicated that there must exist a rapid transport of ice, and perhaps water, across the Arctic Ocean from

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|](#)[|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

runoff was not confined to the coastal areas but extended over the entire Arctic Ocean (Fig. 2 lower panel).

Finally it was found that the salinity increased towards the bottom, forming a high salinity, cold bottom water. Nansen interpretation of these high bottom salinities was that during ice formation over the shallow shelf areas brine was released and sank towards the bottom, where it accumulated during winter gradually increasing in density. Eventually this dense shelf bottom water crossed the shelf break and sank into the deep Arctic Ocean, creating the cold, saline bottom water (Nansen, 1906). The accuracy of the salinity measurements on the Fram salinity samples was later questioned, and after making observations of the Arctic Ocean deep and bottom water north of Svalbard 10 Nansen concluded that the salinities measured on the Fram samples were too high, and proposed that the deep water in the Arctic Ocean derived from the open ocean deep convection area in the Central Greenland Sea (Nansen, 1915).

After the drift of Fram intense oceanographic studies were also conducted in the 15 Greenland Sea and the Barents Sea (e.g., Knipovich, 1905; Nansen, 1906; Helland-Hansen and Nansen, 1909) and the Northwest Passage was finally navigated by Roald Amundsen with Gjøa 1903–1905. Fram under Sverdrup made a second voyage to the Canadian Arctic Archipelago 1898–1902 and explored several of its islands and straits, and the Russian Hydrographic Expedition of the Arctic Ocean 1910–1915 charted the 20 northern sea route (the Northeast Passage) and discovered Severnaya Zemlya. After the first World War Amundsen tried to reach the North Pole from the Pacific on Maud 1918–1925, but Maud was trapped by the ice in the East Siberian Sea and did not cross the shelf break into the deep basin. The studies on tides and long waves by H.U. Sverdrup (1925) and on sea ice by F. Malmgren (1927) were the main oceanographic 25 achievements from this expedition.

In the 1930s the Soviet icebreaker Sedov involuntarily duplicated the drift of Fram, getting caught in the ice. Another, rather spectacular, attempt to penetrate into the interior of the Arctic Ocean was made in 1931, when a discarded US submarine, Nautilus, was taken to the waters west and north of Svalbard. When Nautilus reached the ice, it

Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

was discovered that the diving rudders did not function. The submarine could not dive and no subsurface voyage to the North Pole could be made. It should take 27 years before another Nautilus surfaced at the North Pole. However, some oceanographic stations were taken in the deep Arctic Ocean north of Svalbard. A more successful

attempt to reach the North Pole was made in May 1937, when the Soviet Union established its first ice station North Pole – 1 at 89.4° N, 78.7° E. The station drifted quickly southward and passed through the strait between Greenland and Svalbard, now called Fram Strait after the Fram drift, and the four members were picked up by the Soviet icebreakers Taymyr and Murman in the Greenland Sea in February 1938.

The observations from Nautilus and from North Pole – 1 complemented the picture of the Arctic Ocean given by Fram voyage. It was a deep ocean with a warm Atlantic layer deriving from the inflow through Fram Strait in the West Spitsbergen Current and a cold bottom water with temperatures increasing with increasing distance from the Greenland Sea, supporting Nansen's revised view of the deep water ventilation with the deep water source for the Arctic Ocean and the Nordic Seas in the Greenland Sea (Fig. 3).

After World War II more ice stations were established, by the Soviet Union and by the United States and Canada, and the airborne activity of the Soviet Union continued in spring each year extending over almost the entire Arctic Ocean, gradually increasing the number of hydrographic stations. From these observations a more detailed picture of the bathymetry, the water masses and the circulation of the Arctic Ocean was emerging (Fig. 4). In the following sections we shall concentrate on five themes present almost at the beginning of the scientific study of the Arctic Ocean and see how our understanding of these themes has evolved, if at all, from the first attempts made by Nansen and others in the beginning of the last century. The themes considered are:

1. The low salinity surface layer and the storage and export of freshwater
2. The heat flux between the Atlantic layer, the sea ice and the atmosphere
3. The circulation of the Atlantic water

2 The low salinity surface layer and the storage and export of freshwater

The Arctic Ocean is a small ocean. Its coastline, and hence its drainage area, is large compared to its area. Over the ocean precipitation is larger than evaporation, adding to the freshwater input. This creates the strong stability in the upper layer and allows ice to form in winter and to survive the seasonal melting in summer. Freshwater is also supplied by the low salinity inflow through Bering Strait. This inflow is driven by the higher sea level in the North Pacific compared to the North Atlantic, which ultimately arises because of the atmospheric water vapour transfer from the North Atlantic to the North Pacific across the Isthmus of Panama (Weyl, 1968; Stigebrandt, 1984). However, the transport has a strong seasonal cycle, being weaker in winter due to persisting northerly winds that oppose the inflow (Coachman and Aagaard, 1988). A smaller advective contribution to the positive freshwater balance of the Arctic Ocean is provided by the Norwegian Coastal Current that carries low salinity water from the Baltic and from the Norwegian coast into the Barents Sea and then to the Arctic Ocean via the Siberian shelf seas.

The freshwater balance is maintained by the export of sea ice and by the outflow of low salinity surface water through Fram Strait and through the shallow and narrow passages in the Canadian Arctic Archipelago. The ice export is largely driven by the wind fields and only to a smaller degree by the ocean currents and may therefore be subject to large temporal variations. The transport of liquid freshwater, by contrast, occurs primarily as rotationally controlled buoyant boundary currents. The currents have the coast to the right (in the Northern Hemisphere) and flow above a denser water mass assumed at rest, which reaches the surface away from the coast.

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Following Rudels (2010) we assume a vertical coast and that the transport in the upper layer is given by Werenskiold's formula (Werenskiold, 1935);

$$M = \frac{(\rho_A - \rho_1)gH_1^2}{2\rho_A f} = M_A + F \quad (1)$$

Here H_1 is the depth of the upper layer at the coast, ρ_A and ρ_1 the density of the lower and upper layer respectively, g the acceleration of gravity and f the Coriolis parameter.
5 F is the liquid freshwater export (freshwater input – ice export) and M_A the amount of lower layer water entrained into and exported in the upper layer.

In cold ocean waters the density is almost exclusively determined by the salinity and we can apply a simplified equation of state:

$$10 \quad \rho = \rho_f(1 + \beta S) \quad (2)$$

where ρ_f is the density of freshwater, β the coefficient of salt contraction and S the salinity. The reduced gravity g' can then be written as:

$$g \frac{(\rho_A - \rho_1)}{\rho_A} = g' \approx g\beta(S_A - S_1) \quad (3)$$

where S_A and S_1 are the salinity of the lower (Atlantic) layer and the upper layer and
15 $\rho_f/\rho_A \approx 1$ (e.g., Stigebrandt, 1981).

The export of low salinity surface water then depends upon the number of effective passages, the depth of the upper layer and the salinity difference between the layers in the passages. Of the three passages in the Canadian Arctic Archipelago that connect the Arctic Ocean to Baffin Bay only two, Lancaster Sound and Nares Strait, are wide enough to transport with full capacity (Fig. 4). Assuming that the depth of the upper layer and the salinity of the upper and lower layers are the same in all passages the salinity in the upper layer can be written as:
20

$$S_1 = \frac{M_A S_A}{(F + M_A)} \quad (4)$$

and the freshwater storage m (in metres) in the Arctic Ocean water column becomes:

$$m = \frac{(S_A - S_1)H_1}{S_A} \quad (5)$$

If S_1 is eliminated between Eqs. (4) and (5) we obtain;

$$(M_A + F)^2 = \frac{3gFS_A H_1^2}{2f} \quad (6)$$

5 and

$$m = \frac{FH_1}{(M_A + F)} \quad (7)$$

The freshwater storage can then be written as;

$$m = \left(\frac{2fF}{3g\beta S_A} \right)^{1/2} \quad (8)$$

10 m only depends upon the freshwater input and the salinity in the lower layer, the number of outlets and on f and g . The width of the boundary current is approximately given by the Rossby radius $Ro = \sqrt{gH_1/f}$, which becomes:

$$Ro = \frac{(g\beta(S_A - S_1)H_1)^{1/2}}{f} = \frac{(g\beta S_A m)^{1/2}}{f} = \left(\frac{2g\beta S_A F}{3f^3} \right)^{1/4} \quad (9)$$

and also only depends upon the freshwater storage and ultimately on the freshwater input.

15 Applying reasonably realistic values for the freshwater input, $F = 0.2 \text{ Sv}$ ($1 \text{ Sv} = 1 \times 10^6 \text{ m}^3 \text{ s}^{-1}$), and setting the Atlantic water salinity S_A to 35 and using $g = 9.83 \text{ ms}^{-2}$ and $f = 1.4 \times 10 \text{ s}^{-1}$ the freshwater storage becomes $m = 8.25 \text{ m}$ and the Rossby radius $Ro \approx 10 \text{ km}$ (Rudels, 2010). The freshwater storage estimated by Serreze et al. (2006) was $74,000 \text{ km}^3$, which spread over the entire Arctic Ocean ($10 \times 10^6 \text{ km}^2$) is 7.4 m ,

OSD

8, 2313–2376, 2011

Arctic Ocean
Circulation

B. Rudels

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

fairly close to the value obtained here. A width of the boundary current of 10 km shows that both Lancaster Sound and Nares Strait are wide enough to transport with full capacity.

If the upper layer depth is different in the three channels the outflow will be somewhat larger and the mean depth (and storage) of the freshwater layer in the Arctic Ocean must be somewhat smaller to achieve a freshwater balance. In the situation that there would be no net ice formation and sea ice export, presently estimated to 0.09 Sv, the freshwater input F would rise to 0.29 Sv, and the freshwater storage has to increase to 10.1 m before the export balances the input. In general, the freshwater storage increases as \sqrt{F} , while the residence time decreases as $1/\sqrt{F}$. If only the storage in the Central Arctic Ocean with an area of $0.5 \times 10^6 \text{ km}^2$ is considered, the residence time for the liquid freshwater is presently about 13 yr while the storage time for sea ice with a thickness of 2 m would be around 7 yr.

Much of the exchange through Fram Strait, especially the inflow, takes place in barotropic streams following the bottom contours and through barotropic eddies (Schauer et al., 2008). Although the barotropic transports contribute significantly to the exchanges, perhaps providing the major part of the transports, the barotropic velocities are still considerably smaller than the baroclinic velocities in the upper low salinity water. Adding a barotropic component to the shallow, low salinity surface layer should then not increase, or decrease, the freshwater outflow by any large amount.

Another assumption made is that the density (salinity) of the lower layer is the same in all channels. This is not the case. The salinity of the lower layer in Fram Strait is close to 35 as is used here but the lower layer in the Canadian Arctic Archipelago, represented by the upper layer in Baffin Bay, has a salinity of 33.7 (Riis-Karstensen, 1936). The situation in the Baffin Bay and Canadian Arctic Archipelago is more complicated since there are two, actually three, control sections, two in the Archipelago and one in Davis Strait. This fact can be used to better determine the transports of volume and freshwater west of Greenland (Rudels, 1986a, 2011).

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



layer outflow has been assumed driven only by the baroclinic pressure gradient generated by the freshwater input. The freshwater transport due to a low salinity outflow forced by an imposed sea level difference between the Arctic Ocean and the North Atlantic is not taken into account. An attempt to introduce an additional “barotropic” sea level correction for the Canadian Arctic Archipelago – Baffin Bay system was made by Rudels (2011).

3 The heat flux between the Atlantic layer, the sea ice and the atmosphere

One important effect of the large freshwater supply to the Arctic Ocean is the resulting strong stability that isolates the entering Atlantic water from the atmosphere and insulates the ice cover from the heat transported by the Atlantic Ocean current system from the tropics and subtropics to the Arctic Mediterranean Sea and the Arctic Ocean. This is the effect that prevents the formation of the open Polar Ocean that in the 19th century was speculated to exist. In the 1970s, in the light of a possible diversion of one or two of the Siberian rivers from the Arctic Ocean to Central Asia for irrigation purposes, the same question arose: Will the reduced stability lead to stronger vertical heat flux from the Atlantic layer, thinning and perhaps removing of the ice cover? (Aagaard and Coachman, 1975).

The large transports of Atlantic water into the Arctic Ocean through Fram Strait determined by direct current measurements in the West Spitsbergen Current enhanced the importance of this question (Aagaard et al., 1973; Aagaard and Greisman, 1975). In the early 1980s the Fram Strait Project (FSP) was initiated to determine the exchanges between the Nordic Seas and the Arctic Ocean through Fram Strait. The transports of Atlantic water and heat through Fram Strait were also a focus for the oceanography programme on the Swedish Ymer-80 expedition, one of the first scientific ice breaker expeditions to the Arctic Ocean. CTD sections extending across the entire Fram Strait essentially started with Ymer in 1980 (Rudels, 1987) and were eventually being taken yearly as more powerful research icebreakers like R/V *Polarstern* and

8, 2313–2376, 2011

OSD

Arctic Ocean
Circulation

B. Rudels

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|](#)[|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|](#)[|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

that all lost heat goes to ice melt, be estimated from;

$$S_1 = \frac{S_A}{(1 + cL^{-1}(T_A - T_1))} \quad (10)$$

where S_1 is the salinity and T_1 the temperature of the upper layer and S_A is the salinity of the Atlantic water, $c = 4000 \text{ J kg}^{-1} \text{ degree}^{-1}$ is the heat capacity of seawater and

- ⁵ $L = 335\,000 \text{ J kg}^{-1}$ the latent heat of melting. With $S_A = 35$ and $T_A = 3.1$ and T_1 equal to the freezing temperature -1.9°C giving $(T_A - T_1) = 5^\circ\text{C}$ S_1 becomes ≈ 33 , clearly lower than the observed value.

The higher salinity then indicates that only a fraction, and a minor fraction, goes to ice melt (Fig. 5 Ice melting). The larger part is lost to the atmosphere. (Rudels et al., 1999a) and Rudels (2010) explored the situation, when sea ice melts on warmer

- ¹⁰ water under freezing conditions using a Kato-Phillips energy balance model (Kato and Phillips, 1969). The entrainment velocity w_e then becomes;

$$w_e = \frac{2n_o u_*^3}{g(\beta(S_A - S_1) - \alpha(T_A - T_1))H_1} - \frac{\varepsilon B}{g(\beta(S_A - S_1) - \alpha(T_A - T_1))} \quad (11)$$

Here u_* is the friction velocity, n_o a constant ≈ 1.5 and ε a parameter set to 1 when positive buoyancy is added and to 0.05 when the buoyancy input is negative (Stigebrandt, 1981). α and β are the coefficients of heat expansion and salt contraction, respectively.

As the Atlantic water cools by melting sea ice and by giving up heat to the atmosphere two buoyancy inputs are generated, one positive due to ice melt and one negative due to cooling, and the positive buoyancy input depends on the fraction, ϕ , of heat going to ice melt.

$$B = g \left\{ \left(w_e \frac{\phi c}{L} \Delta T_A \beta S_A + \frac{\phi c}{L} \beta S_A H_1 \frac{d(\Delta T_A)}{dt} \right) - \left(w_e \alpha \Delta T_A + H_1 \alpha \frac{d(\Delta T_A)}{dt} \right) \right\} \quad (12)$$

B is the total buoyancy flux and $\Delta T_A = (T_A - T_1)$. The first bracket is the positive contribution due to ice melt and the second bracket the negative contribution due to cooling.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In the brackets the first terms represent the cooling of the entrained water to the mixed layer temperature and the second terms the reduction of the mixed layer temperature. When the upper layer has reached freezing temperature these two terms are zero. Both buoyancy inputs influence the entrainment velocity, which will be given by:

$$^5 \quad w_e = \frac{n_o u_*^3 (1 + \phi c L^{-1} \Delta T_A)}{g \Delta T_A (\phi c L^{-1} \beta S_A - \alpha (1 + \phi c L^{-1} \Delta T_A)) H_1} - \frac{H_1}{2 \Delta T_A} \frac{d(\Delta T_A)}{dt} \quad (13)$$

The crucial assumption is that the heat flux to ice melt is a minimum (Rudels et al., 1999; Rudels, et al., 1999a). This keeps the density step between the created low salinity surface layer and the Atlantic water below small and allows for a more rapid transfer of heat from the Atlantic water to the upper layer and to ice melt and to the atmosphere. The entrainment velocity is directly related to the heat flux Q and the fraction for which least heat goes to ice melt is found from:

$$Q = G \left\{ u_*^3, \Delta T_A, t, \frac{d(\Delta T_A)}{dt} \right\} \times \frac{1}{\sqrt{\phi c L^{-1} \beta S_A - \alpha (1 + \phi c L^{-1} \Delta T_A)}} \quad (14)$$

where only the last factor depends upon ϕ . The fraction ϕ_o is then determined from;

$$\frac{d(\phi Q')}{d\phi} = 0, \quad \text{with} \quad Q' = \left(\phi c L^{-1} \beta S_A - \alpha (1 + \phi c L^{-1} \Delta T_A) \right)^{-1/2} \quad (15)$$

which gives;

$$\phi_o = \frac{2\alpha L}{c(\beta S_A - \alpha \Delta_A T)} \approx \frac{2\alpha L}{c\beta S_A} \quad (16)$$

The upper layer salinity then becomes;

$$S_1 = \frac{S_A}{\left(1 + \frac{\phi_o c \Delta T_A}{L} \right)} \approx \frac{S_A}{\left(1 + \frac{2\alpha \Delta T_A}{\beta S_A} \right)} \quad (17)$$

Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Using $\alpha = 0.6 \times 10^{-4}$ and $\beta = 8 \times 10^{-4}$ S_1 becomes ~ 34.3 , which should be compared to 33.1 found above for the case when all heat goes to ice melt. This is the minimum salinity. Once the freezing temperature is reached in the upper layer the entrained heat is not sufficient to fulfil the heat flux to the atmosphere and to ice melt and new ice starts to form. This is then considered to occur simultaneously to the ice melt (Rudels et al., 1999a) (Fig. 5).

North of Svalbard the energy driving the entrainment from the Atlantic water to the upper layer is supplied by the wind. As the Atlantic water is advected farther and eastward into the Arctic Ocean in the boundary current along the Eurasian continental slope its upper part becomes stratified by the seasonal ice melt in summer and in winter ice formation and brine rejection generate haline convection. This homogenizes the upper layer down to the thermocline above the Atlantic water. The TS curves that during the wind mixed phase were straight lines from the mixed layer to the temperature maximum of the Atlantic water become curved, indicating that the brine enriched water penetrates into and cools the upper part of the thermocline (Rudels et al., 1996; Kikuchi et al., 2004) (Fig. 6). Convection is not a good stirrer and its intensity depends on the freezing. Should it bring warm water towards the ice the freezing would diminish and the entrainment would cease. The ice cover is thus in a sense self protected against convectively driven heat flux from below (Martinson and Steele, 2001).

The second inflow branch from the Nordic Seas to the Arctic Ocean passes over the Barents Sea. It is cooled extensively in the southwestern part of the Barents Sea and also becomes freshened by net precipitation. The temperature and salinity of the core of the Atlantic water become lower than those of the Fram Strait inflow before it encounters sea ice. This should, according to Eq. (17), lead to less heat going to ice melt and more heat going to the atmosphere. The salinity of the upper, transformed, layer then becomes higher in the Barents Sea branch than in the Fram Strait branch. The two inflows meet north of the Kara Sea as the main part of the Barents Sea inflow enters the Arctic Ocean in the St. Anna Trough. On a north-south section taken over the shelf and slope by Polarstern in 2007 crossing the two inflow branches the TS

Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

curves from the two branches clearly show that the upper part of the Barents Sea branch has a higher salinity than the warm Fram Strait branch (Fig. 7).

It is also evident that the salinity at the base of the upper layer, which indicates the salinity of the mixed layer in winter, in both branches better corresponds to the salinity obtained from Eq. (17) than the situation when all heat goes to ice melt (Eq. 10). The lower salinity seen along the freezing point line reflects the summer melting caused by the incoming short wave solar radiation.

North of the Laptev Sea the boundary current and the two branches become overrun by low salinity shelf water comprising the major part of the river runoff from Ob, Yenisey and Lena that crosses the shelf break. Cooling and ice formation are not strong enough to homogenize the water column down to the thermocline but only to the base of the less saline shelf water layer that becomes the Polar Mixed Layer (PML). The upper parts of the two inflow branches then become isolated from the surface processes and are transformed to halocline waters that further insulate the sea ice cover from the heat stored in the Atlantic layer. The mixing and entrainment that do occur will only bring more saline but cold halocline water into the mixed layer and little heat is supplied from below. The heat from the Atlantic layer is instead trapped in the halocline, where the temperature gradually increases (Fig. 6).

The winter convection and the homogenization of the PML is in summer identified by an upper temperature minimum that is the remnant of the convection being eroded by the radiative heating above and by mixing with the somewhat warmer halocline below, where the heat lost vertically upward by the Atlantic layer is stored (Fig. 8). Only on the slope in the Western Nansen Basin, where no halocline is present and the Atlantic layer is brought close to the sea surface, does the temperature minimum become completely eroded. This indicates that heat can be transferred from the Atlantic layer to the mixed layer and the ice also in summer. In the rest of the Arctic Ocean the heat transferred upward from the Atlantic layer will be trapped in the halocline (Rudels, 2010).

The heat flux appears to be stronger in the Barents Sea branch, probably due to enhanced mixing over the continental slope. The colder Fram Strait branch is farther

away from the slope and eventually supply the halocline in the Amundsen, Makarov and Northern Canada Basins, while the Barents Sea branch remains at the slope and passes between the Chukchi Sea and the Chukchi Cap into the Southern Canada Basin (Rudels et al., 2004; Shimada et al., 2005).

- 5 Between the Laptev Sea and the Canada Basin shelf water is added to the Arctic Ocean as well as Pacific water from Bering Strait via the Chukchi Sea. Most of these waters are less dense than the Laptev Sea outflow and contribute further layers to the upper part of the water column. Some water masses like the Bering Sea Summer Water BSSW and Bering Sea Winter Water (BSWW) (Coachman and Barnes, 1961) and
10 other brine enriched shelf water from the Chukchi Sea (Jones and Anderson, 1986) are dense enough to penetrate between the PML and the halocline, making further contributions to the halocline and making the initial, denser halocline with salinity of ~34.2 into the lower halocline (Fig. 8). The circulation of the halocline waters is summarized in Fig. 9.

15 4 The circulation of the Atlantic water

The uppermost part of the water column follows the wind driven ice motion and circulates anticyclonally in the Beaufort Gyre in the Southern Canada Basin and also participates in the transpolar drift from Siberia towards Fram Strait. By contrast the
20 Atlantic water moves cyclonically around the deep Arctic Ocean Basin. This motion was first deduced from the decrease in temperature of the Atlantic core in the different parts of the Arctic Ocean. It is highest in the Nansen Basin north of Svalbard and lowest in the Canada Basin and north of Greenland, indicating a cyclonic flow and a gradual cooling of the Atlantic layer (Timofeyev, 1960; Coachman and Barnes, 1963).

In 1991 the icebreaker Oden took a hydrographic section from the Central Nansen Basin across the Gakkel Ridge, the Amundsen Basin, the Lomonosov Ridge into the Makarov Basin (Anderson et al., 1994; Rudels et al., 1994). On the section two distinct temperature fronts were found, one over the Gakkel Ridge and one over the Lomonosov Ridge. At the same time extensive layers of temperature and salinity

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



inversions were observed. The layers, being identified in TS space on several stations, suggested mixing between two water masses, one warm and saline closer to the continental slope and the other colder and fresher closer to the Lomonosov Ridge. The intrusive layers were found over a large depth range and the transports of the two involved water masses should be of about equal strength. The origin of the warmer end member is the inflow through Fram Strait and the most likely source for the colder end member is the inflow from the Norwegian Sea over the Barents Sea to the Arctic Ocean. The Barents Sea branch enters the Arctic Ocean in the St. Anna Trough and should be found closer to the continental slope north of the Kara and Laptev seas.

The Oden section did not cover the continental slope east of the St. Anna Trough and the properties of this inflow had to be hypothetical. Furthermore, since the intrusions were observed in the central part of the basins with the warmest water in the Nansen Basin and the colder, less saline water in the Amundsen Basin, this would imply that a part of the boundary current leaves the continental slope and enters the interior Eurasian Basin, forming a return flow towards Fram Strait. The intrusions did not seem to change substantially between the main Oden section and the sections taken closer to Fram Strait, indicating that they were no longer expanding laterally but were advected as fossil structures with the main flow. At the Lomonosov Ridge the Atlantic core was somewhat warmer and this was tentatively explained by a more rapid return flow along the Lomonosov Ridge (Rudels et al., 1994).

There was also a strong contrast between the Atlantic and intermediate waters on the two sides of the Lomonosov Ridge. The Atlantic water was much colder and less saline, while the intermediate water was warmer and more saline in the Makarov Basin than in the Amundsen Basin. This suggested that water on the Makarov Basin side was moving from Greenland towards Siberia after it had travelled around the Makarov Basin. The change in water mass properties would then be due to shelf-slope convection, where initially cold, saline plumes, created by brine rejection on the shelves, were sinking down the slope, entering and cooling the Atlantic water and also bypassing the Atlantic layer entraining warmer water and displacing it to deeper levels. Following

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

a scheme proposed by Rudels (1986b) a plume entrainment model was applied to described the changes in the Atlantic and intermediate water in the Amerasian Basin and to determine the amount and the properties of the shelf water needed to explain the observed changes in the water column (Rudels et al., 1994) (Fig. 10).

5 The postulated colder and less saline inflow from the Barents Sea via the St. Anna Trough was observed north of the Laptev Sea in 1993 (Schauer et al., 1997) and intense interactions between the two inflow branches were seen on the slope north of Severnaya Zemlya in 1995 (Rudels et al., 2000). Nutrient observations from the Atlantic and intermediate waters north of Greenland also suggested that the coldest
10 Atlantic water closest to Greenland actually represented a separate stream that had passed along the continental slope through the Canada Basin and was returning along the North American and Greenland continental slope to Fram Strait (Rudels et al., 1994).

In the 1990s the first reports and studies of changes in the Atlantic water temperature
15 were published (Quadfasel et al., 1991; Carmack et al., 1995; Morison et al., 1998) and subsequent expeditions and studies could follow the propagation of the warm inflow pulse. Its movement largely confirmed the proposed circulation scheme for the Atlantic water (Fig. 11). Some additional loops were found (Smethie et al., 1998; Smith et al., 1999) but also the spreading of the warmer water into the Southern Canada Basin
20 appeared uncertain. Did it enter the Canada Basin between the Chukchi Cap and the Chukchi Continental slope or did it moved around the Chukchi Cap (Shimada et al., 2004)? Was the heat transported into the Canada Basin as thermohaline intrusions (Walsh and Carmack, 2003; McLaughlin et al., 2009) and did it circulate anticyclonally around the Beaufort Gyre (McLaughlin et al., 2009) as was previously suggested by
25 Coachman and Barnes (1963) and Newton and Coachman (1974). Coachman and Barnes (1963) and Newton and Coachman (1974), however, report that a cyclonal circulation along the continental slope appeared to be present also in the Canada Basin (Fig. 11). This then makes the name “the Arctic Circumpolar Boundary Current” used by Rudels et al. (1999b) still valid. It also became clear that much of the cooling of

the Atlantic water and the warming of the intermediate water must occur north of the Chukchi shelf, since the water on the East Siberian Sea shelf had too low salinity to produce water dense enough to sink beneath the PML (Aagaard et al., 1981).

Several warm Atlantic water pulses have later been observed and followed upstream in the Norwegian Sea and downstream in the boundary current along the Eurasian continental slope, along the Lomonosov Ridge and into the Amundsen Basin (Swift et al., 1997; Polyakov et al., 2005). The suggested return flow in the northern Nansen Basin and over the Gakkel Ridge has, however, largely been questioned or ignored.

The area north of the Laptev Sea is where the largest changes in the Atlantic water properties take place. The core of the Atlantic water does not appear to cool significantly in the boundary current between the western Barents Sea (30° E) and the eastern Kara Sea (90° E) and smooth profiles in temperature and salinity are seen in the core of the Fram Strait branch (Fig. 12). In the Kara Sea the inner part of the boundary current comprises the Barents Sea inflow branch that is located at the 400 m isobath and the potential for mixing between the two branches is present and partly takes place. North of the Laptev Sea the smooth central core of the Fram Strait branch has disappeared. The temperature of the temperature maximum has decreased by 1°C and the salinity has been reduced by 0.05. Strong intrusions are also present in the Atlantic layer and in the intermediate water below. The temperature is lower at the slope and in the interior of the basin and lateral temperature and salinity maxima are still present close to the slope (Fig. 13). At 90° E, on the basin side of the Fram Strait branch, similar cooling and freshening of the Atlantic and intermediate waters are observed and intrusions indicate lateral mixing with the less saline and colder Barents Sea branch (Fig. 12).

There are four processes that can explain the strong cooling and freshening of the Atlantic layer that occur north of the Laptev Sea. (1) Heat loss to the atmosphere and to ice melt. The difficulty with this explanation is to introduce freshwater into the core of the Atlantic layer since melt water, due to its low density, would likely be trapped in the Polar Mixed Layer. (2) Lateral mixing between the two inflow branches, leading to

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean Circulation

B. Rudels

the formation of intrusive layering and to lower temperature and salinity in the Fram Strait branch. The heat remains in the core but the temperature is reduced because of the mixing and volume increase due to the incorporation of Barents Sea branch water. (3) Cooling and freshening by shelf-slope convection that brings colder, less saline water into the Atlantic layer and also convects denser water that bypasses the Atlantic layer, entrains warmer water and brings it downwards to deeper levels. Heat is displaced vertically but stays in the water column. (4) The Fram Strait branch is imbedded between Barents Sea branch water at the slope and in the interior, which suggests that the Barents Sea branch water partly leaves the slope and enters the interior of the Nansen Basin. The Fram Strait branch would then have to return with the Barents Sea branch water towards Fram Strait within the Nansen Basin and most of its heat would not pass eastward of the section along 126° E and thus not enter the Amundsen Basin and cross the Lomonosov Ridge. This return flow then also carries the intrusive layers between the two branches towards Fram Strait.

This implies that most of the Atlantic layer in the Arctic Ocean beyond the Nansen Basin is supplied by the Barents Sea branch. By comparing the TS structure of the different basins with that of the Barents Sea branch, this does not appear inconceivable (Fig. 14). However, there are no watertight boundaries and intermittently some water from the Fram Strait branch enters the Barents Sea branch and crosses the Lomonosov Ridge. The warm pulse traced since the 1990s could be such an instance.

Another caveat is that the Barents Sea branch properties considered here are those observed on the shelf-slope at 90° E. This may not be the properties that the Barents Sea branch had, when it left the Barents Sea and entered the St. Anna Trough. Recent studies indicate that most heat of the Atlantic water entering the Barents Sea is lost locally and the average temperature of the Atlantic water leaving the Barents Sea is about 0 °C (Gammelsrød et al., 2010). However, most of the surface water is at the freezing point and the deeper layers have temperatures between –1.0 °C and –0.5 °C. A remaining core of Atlantic water with temperatures around 1 °C could be present and still give an average outflow temperature of 0 °C.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	

Printer-friendly Version
Interactive Discussion



It is also possible that the Fram Strait branch water that enters the St. Anna Trough from the Arctic Ocean mixes with the Barents Sea branch in the trough and becomes incorporated in the Barents Sea branch, adding warmer Atlantic water, and then flows high on the slope eastward within the Barents Sea branch. In such a way the Fram
5 Strait branch would contribute to the Atlantic layer also beyond the Nansen Basin.

5 The deep and bottom waters

Differences between in the water columns in the basins indicate that water mass transformations are taking place, changing the properties of the water masses as they are advected through the Arctic Mediterranean Sea. Because of the strong stability in the

10 upper part of the Arctic Ocean any changes occurring in the deeper layers must be caused by processes taking place at the rim. The most obvious mechanism for creating dense water capable of sinking into the deep basins is, as Nansen (1906) already pointed out, freezing and brine rejection over shallow areas, where the brine can accumulate and generate cold, highly dense bottom water. Evidence that such process is
15 active had already been found e.g. close to the coast of Novaya Zemlya, where a lee polynya forces ice away from the coast, allowing for larger ice production and stronger brine rejection (Knipovich, 1905) (Fig. 15).

As the cold, saline waters cross the shelf break and sink down the slope into the deep basins they entrain ambient water and become less dense. Depending upon their
20 initial density they sink to different depths, where the final density matches that of the surroundings. There they merge with and change the characteristics of the boundary current. Initially less dense plumes supply the halocline in the Canada Basin (Jones and Anderson, 1986) and cool and freshen the Atlantic layer, while denser plumes sink deeper, entraining Atlantic water and become warmer than the ambient waters and
25 thus both heat and increase the salinity of the deeper layers (Rudels, 1986b) (Fig. 16). The characteristics of the Atlantic, intermediate and deep waters in the different basins largely agree with such a process. Compared with the Greenland Sea, the second

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



6 The exchanges through Fram Strait

Fram Strait is the deepest (2600 m) and the most important connection between the Arctic Ocean and the World Ocean. It has a two directional exchange. To the east warm Atlantic water and also Arctic intermediate and deep waters from the Nordic Seas are carried northward in the West Spitsbergen Current, while in the west the East Greenland Current transports low salinity Polar surface water, cooled and transformed Arctic Atlantic Water and different Arctic Ocean intermediate and deep waters southward to the Nordic Sea (Fig. 20). The net transport is about 2 Sv out of the Arctic Ocean and the total in- and outflows are around 7–9 Sv and 9–11 Sv, respectively (Schauer et al., 2008; Rudels et al., 2008). The flow has a strong barotropic component, strongest in the West Spitsbergen Current but significant also in the East Greenland Current, which makes geostrophic calculations unreliable and subjective, depending upon how the unknown barotropic component is handled or approximated. Direct current measurements have been carried out in Fram Strait since 1997 by the Alfred Wegener Institute and the Norwegian Polar Institute with a mooring array consisting of 16 individual moorings. Here difficulties also arise due to the presence of baroclinic and barotropic eddies with small horizontal scales, which complicate the transport estimates.

Perhaps a more fundamental question is – how are the exchanges through Fram Strait driven? – are they mainly forced by the atmospheric circulation or are the thermodynamic processes taking place in the Arctic Ocean and in the Nordic Seas of equal, or greater, importance? To make an extreme case we ask the question: is it possible to have a geostrophically balanced exchange through Fram Strait so strong that the transformations occurring in the Arctic Ocean are just sufficient to create the density field in Fram Strait required for fulfilling the transports needed for the transformations?

To examine this question we construct two water columns, one representing the West Spitsbergen Current the other the East Greenland Current. The geostrophic flow is driven by the density differences between the two water columns and the sea level slope. Following Rudels (1989) we further assume that there is no flow against the

OSD

8, 2313–2376, 2011

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

negative pressure gradient. This means that there will be a southward flow in the upper part of the East Greenland Current, while there is no northward transport in the West Spitsbergen Current at the same level. First below the pressure reversal, where the negative pressure gradient is directed into the Arctic Ocean, or from the 5 West Spitsbergen Current to the East Greenland Current, does the Atlantic water flow northward into the Arctic Ocean. At this level the outflow in the East Greenland Current is blocked.

We only consider Fram Strait. The inflow through Bering Strait is taken to exit through the Canadian Arctic Archipelago and the inflow over the Barents Sea is presently ignored. The river runoff is assumed first to be frozen to sea ice and then a part of the 10 sea ice is melted by the heat supplied by the inflowing Atlantic water. Taking the Atlantic water temperature to be 3.1°C and the salinity 35 and using the approximation of the density, ρ , given by;

$$\rho = \rho_f(1 - \alpha(T - 0.6) + \beta S) \quad (18)$$

15 Where 0.6°C is the mean temperature of the Atlantic water set to 3.1°C and the outflow at the freezing point, -1.9°C . $\alpha = 0.6 \times 10^{-4}$ and $\beta = 8 \times 10^{-4}$ (Gill, 1982). From the expression for the salinity of the upper layer, Eq. (17), the upper layer salinity, S_1 , becomes 34.27. A possible salinity increase due to brine rejection is ignored.

The river runoff to the Arctic Ocean, and thus the ice export, is about 0.1 Sv and we 20 first assume that 0.02 Sv is melted by heat carried by the Atlantic water. The amount of low salinity Polar water leaving the Arctic Ocean can then be computed from the Knudsen relations (Knudsen, 1900):

$$M_1 = M_A + 0.2, \quad (19)$$

$$S_1 M_1 = S_A M_A \quad (20)$$

25 The depth of the upper layer is then found from Werenskiold's formula (Eq. 1) with H_1 being the depth of the pressure reversal given by:

$$\rho_1 H_1 = \rho_A (H_1 - \delta) \quad (21)$$

Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

where δ is the sea level difference between the two water columns. The inflow of Atlantic water takes place below the upper layer depth H_1 , which is also the depth of the pressure reversal, and to transport the needed volume M_A a thickness H_A is required, which can be determined from the geostrophic transport;

$$M_A = \frac{g}{\rho f} \left\{ (\rho_A - \rho_1) \frac{H_A}{2} \right\} H_A \quad (22)$$

However, the pressure gradient driving the Atlantic water cannot be removed with depth even if the depth of the upper layer in the East Greenland Current would be less than $H_1 + H_A$, and the Atlantic water will continue to flow into the Arctic Ocean in the deep (Fig. 21). This leads to an imbalance of the exchanges and the sea surface in the Arctic Ocean would start to rise. A barotropic outflow is then generated, which balances the inflow. The exchanges become larger than those just necessary to create the required density field.

So far the fact that also denser waters are produced in the Arctic Ocean has not been taken into account. To simplistically do so we allow some of the low salinity Polar water to flow onto the shelves, where its salinity is increased by brine rejection to 35 while its temperature remain at the freezing point, -1.9 . In the three examples considered, 25 %, 50 % and 75 % of the Polar surface water is assumed to flow onto the shelves (Tables 1 and 2). As the shelf water descends down the slope, it entrains ambient water and we consider five cases, where the entrainment increases the volume of the convecting shelf water by 0 %, 20 %, 40 %, 60 % and 80 % (Tables 1 and 2). The salinity 35 for layer 3 is chosen to simplify the density computations of the outflowing deep water, which then has the same salinity as the Atlantic water and only differs in temperature. The density of the deep water should be considered as an average of the denser waters formed in the Arctic Ocean, the Arctic Atlantic water and the intermediate and the deep waters.

To export the deep water from the Arctic Ocean a second pressure reversal must exist, where the negative pressure gradient is directed out of the Arctic Ocean. The

Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

second pressure reversal is given by;

$$g\rho_A(H' + H'') = g\rho_1H' + g\rho_3H'' \text{ or } g(\rho_A - \rho_1)H' = -g(\rho_A - \rho_3)H'' \quad (23)$$

where H' is the thickness of the Atlantic inflow layer that lies beside the upper Polar water and the pressure gradient increases with depth, and H'' is the thickness of the part of the Atlantic layer that lies beside the deep water and the pressure gradient decreases with depth. The transport M_A in the Atlantic layer than becomes (Rudels, 1989);

$$M_A = \frac{g}{\rho f} \left\{ \left[(\rho_A - \rho_1) \frac{H'}{2} \right] H' + \left[(\rho_A - \rho_1)H' + (\rho_A - \rho_3) \frac{H''}{2} \right] H'' \right\} \quad (24)$$

Since M_A is known from the transformations, H' and H'' , can be determined from Eqs. (23) and (24). The minimum thickness of the outflow of deep water is then found by solving;

$$M_3 = \frac{g}{\rho f} \left\{ (\rho_3 - \rho_A) \frac{H_3}{2} \right\} H_3 \quad (25)$$

for H_3 , where H_3 is the layer thickness below the second pressure reversal where the export of dense water M_3 takes place (Table 2). However, again similar difficulties arise. The pressure gradient will continue to increase and the deep southward flow will continue below the depth, where mass balance is achieved ($H_1 + H' + H'' + H_3$) (Fig. 22). The sea level in the Arctic Ocean will decrease and generate a compensating barotropic inflow to maintain mass balance.

It thus appears that it is not possible to create a baroclinic exchange that exactly balances the changes in water mass properties. The water mass transformations generate a barotropic flow component through the strait. This would then add to any wind driven barotropic transport created by the atmospheric circulation.

These simple considerations allow for the construction of temperature and salinity fields as well as the current field in Fram Strait for different estimates of Atlantic water

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



remain fairly constant because of the large residence time in the Arctic Ocean (> 20 yr). The recirculation as well as the barotropic and baroclinic eddies present in the central part of Fram Strait are not considered at all (Fig. 23). This is largest part of the strait and the assumption that these transports carry water into and out of the Arctic Ocean but do not contribute significantly to the water mass transformation in the Arctic Ocean has to be demonstrated.

7 Summary

This overview has presented some instances in the history of the exploration and study of the Arctic Ocean, and some basic themes in Arctic Ocean oceanography

have been examined, following ideas and using methods introduced by oceanographers in the early 20th century (Fig. 24): the vision, exploration and insights of Nansen (1902, 1906), Knudsen's relations (Knudsen, 1900), the dynamic method introduced in oceanography by Sandström and Helland-Hansen (1903) and TS analysis first employed by Helland-Hansen (1916). Concepts and ideas that focus on, and are essential for, understanding the water mass transformations and transports taking place in the Arctic Ocean. One concept has largely been neglected here, the wind driven ocean circulation. This is perhaps the most important part for describing the circulation, if not the water mass transformations, in the Arctic Ocean and was also first formulated by Ekman (1905) after observations from Nansen's voyage with Fram.

Acknowledgement. Financial support for this work has been given by the EU project THOR (ThermoHaline Overturning – at Risk?) contract number 212643. Rainer Schlitzer's Ocean Data View has been used to make many of the figures. The historical information given in Sect. 1 has mainly been taken from Laktionov (1960) and Heyes (2003).

OSD

8, 2313–2376, 2011

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



References

Arctic Ocean Circulation

B. Rudels

- Aagaard, K.: On the deep circulation in the Arctic Ocean, *Deep-Sea Res.*, 27, 251–268, 1980.
- Aagaard, K. and Coachman, L. K.: Toward an ice-free Arctic Ocean, *EOS T. Am. Geophys. Un.*, 56, 484–486, 1975.
- 5 Aagaard, K. and Greisman, P.: Towards a new mass and heat budget for the Arctic Ocean, *J. Geophys. Res.*, 80, 3821–3827, 1975.
- Aagaard, K., Darnall, C., and Greisman, P.: Year-long measurements in the Greenland-Spitsbergen passage, *Deep-Sea Res.*, 20, 743–746, 1973.
- 10 Aagaard, K., Coachman, L. K., and Carmack, E. C.: On the halocline of the Arctic Ocean, *Deep-Sea Res.* 28, 529–545, 1981.
- Aagaard, K., Swift, J. H., and Carmack, E. C.: Thermohaline circulation in the Arctic Mediterranean Seas, *J. Geophys. Res.*, 90, 4833–4846, 1985.
- Anderson, L. G., Björk, G., Holby, O., Jones, E. P., Kattner, G., Koltermann, K.-P., Liljeblad, B., Lindegren, R., Rudels, B., and Swift, J. H.: Water masses and circulation in the Eurasian Basin: results from the Oden 91 Expedition, *J. Geophys. Res.*, 99, 3273–3283, 1994.
- 15 Björk, G. and Winsor, P.: The deep waters of the Eurasian Basin, Arctic Ocean: geothermal heat flow, mixing and renewal, *Deep-Sea Res. Pt. I*, 53, 1253–1271, 2006
- Björk, G., Jakobsson, M., Rudels, B., Swift, J. H., Anderson, L. G., Darby, D. A., Backman, J., Coakley, B., Winsor, P., Polyak, L., and Edwards, M.: Bathymetry and deep-water exchange across the central Lomonosov Ridge at 88°–89° N, *Deep-Sea Res. Pt. I*, 54, 1197–1208 doi:10.1016/j.dsr.2007.05.010, 2007.
- 20 Carmack, E. C., Macdonald, R. W., Perkin, R. G., and McLaughlin, F. A.: Evidence for warming of Atlantic water in the Southern Canadian Basin, *Geophys. Res. Lett.*, 22, 1961–1964, 1995.
- 25 Carmack, E. C., Aagaard, K., Swift, J. H., Macdonald, R. W., McLaughlin, F. A., Jones, E. P., Perkin, R. G., Smith, J. N., Ellis, K. M., and Killius, L. R.: Changes in temperature and tracer distributions within the Arctic Ocean: results from the 1994 Arctic Ocean section, *Deep-Sea Res. Pt. II*, 44, 1487–1502, 1997.
- Coachman, L. K. and Aagaard, K.: Transports through Bering Strait: annual and interannual variability, *J. Geophys. Res.*, 93, 15535–15539, 1988.
- 30 Coachman, L. K. and Barnes, C. A.: The contribution of Bering Sea water to the Arctic Ocean, *Arctic*, 14, 147–161, 1961.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Coachman, L. K. and Barnes, C. A.: The movement of Atlantic water in the Arctic Ocean, *Arctic*, 16, 8–16, 1963.
- Ekman, V. W.: On the influence of earth's rotation on ocean currents, *Arch. Math. Astron. Phys., Arkiv för matematik, astronomi och fysik*, 2(11) 1–53, 1905.
- 5 Foldvik, A., Aagaard, K., and Törresen, T.: On the velocity field of the East Greenland Current, *Deep-Sea Res.*, 35, 1335–1354, 1988.
- Gammelsrød, T., Leikvin, Ø., Lien, V., Budgell, W. P., Loeng, H., and Maslowski, W.: Mass and heat transports in the NE Barents Sea: observations and models, *J. Marine Syst.*, 75, 56–69, 2009.
- 10 Helland-Hansen, B.: Nogen Hydrografiske Metoder, *Forh. Skandinaviske Naturf. Mote*, Kristiania, pp. 357, 1916.
- Helland-Hansen, B. and Nansen, F.: The Norwegian Sea, Ist physical oceanography based upon the Norwegian researches 1900–1904, *Rep. on Norw. Fishery and Marine Investigations II(1)*, Kristiania, 1909.
- 15 Heyes, D.: Historical Atlas of the Arctic, Univ, Wash. Press, Seattle, 208 pp., 2003.
- Jakobsson, M., Macnab, R., Mayer, L., Anderson, R., Edwards, M., Hatzky, J., Schenke, H. W., and Johnson, P.: An improved bathymetric portrayal of the Arctic Ocean: implications for ocean modelling and geological, geophysical and oceanographic analyses, *Geophys. Res. Lett.*, 35, L07602, doi:10.1029/2008GL033520, 2008.
- 20 Jones, E. P. and Anderson, L. G.: On the origin of the chemical properties of the Arctic Ocean halocline, *J. Geophys. Res.*, 91, 10759–10767, 1986.
- Jones, E. P., Rudels, B., and Anderson, L. G.: Deep waters of the Arctic Ocean: origins and circulation, *Deep-Sea Res.*, 42, 737–760, 1995.
- 25 Kikuchi, T., Hatakeyama, K., and Morison, J. H.: Distribution of convective lower halocline water in the Eastern Arctic Ocean, *J. Geophys. Res.*, 109, C120301, doi:10.1029/2003JC002223, 2004.
- Knipowitsch, N.: Hydrologische Untersuchungen im Europäischen Eismeer, *Annalen der Hydrographie und Maritimen Meteorologie*, Sonderabdruck, 62 pp., 1905.
- 30 Knudsen M.: Ein hydrographischer Lehrsatz, *Annalen der Hydrographie und Maritimen Meteorologie*, 28, 316–320, 1900.
- Laktionov, A.: Nordpolen, *Förlaget för litteratur på främmande språk*, Moskva, 522 pp, 1960.
- Martinson, D. G. and Steele, M.: Future of the Arctic Sea Ice Cover: Implications of an Antarctic Analog, *Geophys. Res. Lett.*, 28(2), 307–310, 2001.

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

Abstract	Introduction
Conclusions	References
Tables	Figures

[|◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- McLaughlin, F. A., Carmack, E. C., Macdonald, R. W., and Bishop, J. K. B.: Physical and geochemical properties across the Atlantic/Pacific water mass boundary in the Southern Canada Basin, *J. Geophys. Res.*, 101, 1183–1197, 1996.
- 5 McLaughlin, F. A., Carmack, E. C., MacDonald, R. W., Weaver, A. J., and Smith, J.: The Canada Basin 1989–1995: upstream events and far-field effects of the Barents Sea branch, *J. Geophys. Res.*, 107(C7), 3082, doi:10.1029/2001JC000904, 2002.
- 10 McLaughlin, F. A., Carmack, E. C., Williams, W. J., Zimmermann, S., Shimada, K., and Itoh, M.: Joint effects of boundary currents and thermohaline intrusions on the warming of Atlantic water in the Canada Basin, 1993–2007, *J. Geophys. Res.*, 114, C00A12, doi:10.1029/2008JC005001, 2009.
- Morison, J. H., Steele, M., and Anderson, R.: Hydrography of the upper Arctic Ocean measured from the nuclear submarine USS Pargo, *Deep-Sea Res. Pt. I*, 45, 15–38, 1998.
- Nansen, F.: Oceanography of the North Polar Basin, *The Norwegian North Polar Expedition 1893–1896, Scientific Results* (9), 427 pp., 1902.
- 15 Nansen, F.: Northern Waters, Captain Roald Amundsen's oceanographic observations in the Arctic Seas in 1901, *Videnskab-Selskabets Skrifter 1. Matematisk-Naturvidenskabelig Klasse 1, Kristiania*, 1–145, 1906.
- Nansen, F.: Spitsbergen Waters, *Videnskabs-selskabets skrifter I. Matematisk-Naturvidenskabelig klasse I(3)*, 145 pp., 1915.
- 20 Petermann, A.: Der Nordpol und Südpol, die Wichtigkeit ihrer Erforschung in geographischer und kulturhistorischer Beziehung, mit Bemerkungen über die Strömungen der Polar-Meere, *Pet. Mitt.*, Gotha, 146–160, 1865.
- Rabe, B., Karcher, M., Schauer, U., Toole, J. M., Krichfield, R. A., Pisarev, S., Kauker, F., Gerdes, R., and Kikuchi, T.: An assessment of Arctic Ocean freshwater content changes from 1990s to the 2006–2008 period, *Deep-Sea Res. Pt. I*, 58, 175–185, doi:10.1016/j.dsr.2010.12.002, 2010.
- 25 Riis-Carstensen, E.: The “Godthaab” Expedition 1928, *The Hydrographic Work and Material, Meddeleser om Grönland* 78(3), 101 pp., 1936.
- Rudels, B.: The outflow of Polar Water through the Arctic Archipelago and the oceanographic conditions in Baffin Bay, *Polar Res.*, 4 n.s., 161–180, 1986a.
- 30 Rudels, B.: The Θ-S relations in the northern seas: implications for the deep circulation, *Polar Res.*, 4 n.s., 133–159, 1986b.
- Rudels, B.: On the mass balance of the Polar Ocean, with special emphasis on the Fram Strait,

Arctic Ocean Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Rudels, B.: The formation of Polar Surface Water, the ice export and the exchanges through the Fram Strait, *Prog. Oceanogr.*, 22, 205–248, 1989.

Rudels, B.: The thermohaline circulation of the Arctic Ocean and the Greenland Sea, *Philos. T. R. Soc. Lond. A*, 352, 287–299, 1995.

Rudels, B.: Arctic Ocean circulation, In: *Encyclopedia of Ocean Sciences*, edited by: Steele, J. H., Turekian, K. K., and Thorpe, S. A., Academic Press, San Diego, 211–225, doi:016/B978-012374473-9.00601-9, 2009.

Rudels, B.: Constraints on exchanges in the Arctic Mediterranean – do they exist and can they be of use? *Tellus A*, 62, 109–122, 2010.

Rudels, B.: Volume and freshwater transports through the Canadian Arctic Archipelago – Baffin Bay system, 116, C00D10, doi:i10.1029/2011JC007019, 2011.

Rudels, B. and Quadfasel, D.: Convection and deep water formation in the Arctic Ocean – Greenland Sea system, *J. Marine Syst.*, 2, 435–450, 1991.

Rudels, B., Jones, E. P., Anderson, L. G., and Kattner, G.: On the intermediate depth waters of the Arctic Ocean, in: *The Role of the Polar Oceans in Shaping the Global Climate*, edited by: Johannessen, O. M., Muench, R. D., and Overland, J. E., American Geophysical Union, Washington D.C., 33–46, 1994.

Rudels, B., Anderson, L. G., and Jones, E. P.: Formation and evolution of the surface mixed layer and the halocline of the Arctic Ocean, *J. Geophys. Res.*, 101, 8807–8821, 1996.

Rudels, B., Friedrich, H. J., Hainbucher, D., and Lohmann, G.: On the parameterisation of oceanic sensible heat loss to the atmosphere and to ice in an ice-covered mixed layer in winter, *Deep-Sea Res. Pt. II*, 46, 1385–1425, 1999a.

Rudels, B., Friedrich, H. J., and Quadfasel, D.: The Arctic circumpolar boundary current, *Deep-Sea Res. Pt. II*, 46, 1023–1062, 1999b.

Rudels, B., Muench, R. D., Gunn, J., Schauer, U., and Friedrich, H. J.: Evolution of the Arctic Ocean boundary current north of the Siberian shelves, *J. Marine Syst.*, 25, 77–99, 2000.

Rudels, B., Jones, E. P., Schauer, U., and Eriksson, P.: Atlantic sources of the Arctic Ocean surface and halocline waters, *Polar Res.*, 23, 181–208, 2004.

Rudels, B., Marnela, M., and Eriksson, P.: Constraints on estimating mass, heat and freshwater transport in the Arctic Ocean: an exercise, in: *Arctic-Subarctic Ocean Fluxes*, edited by: Dickson, R. R., Meincke, J., and Rhines, P., Springer, Dordrecht, 315–341, 2008.

Sandström, J. W. and Helland-Hansen, B.: Über die Berechnung von Meereströmungen, Rep.

Arctic Ocean Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Schauer, U., Muench, R. D., Rudels, B., and Timokhov, L.: Impact of Eastern Arctic Shelf water on the Nansen Basin intermediate layers, *J. Geophys. Res.*, 102, 3371–3382, 1997.

5 Schauer, U., Beszczynska-Möller, A., Walczowski, W., Fahrbach, E., Piechura, J., and Hansen, E.: Variation of measured heat flow through the Fram Strait between 1997–2006, in: Arctic-Subarctic Ocean Fluxes, edited by: Dickson, R. R., Meincke, J., and Rhines, P., Springer, Dordrecht, 65–85, 2008.

10 Serreze, M. C., Barrett, A., Slater, A. J., Woodgate, R. A., Aagaard, K., Steele, M., Moritz, R., Meredith, M., and Lee, C.: The large-scale freshwater cycle in the Arctic, *J. Geophys. Res.*, 111, C11010, doi:10.1029/2005JC003424, 2006.

Shimada, K., McLaughlin, F. A., Carmack, E. C., Proshutinsky, A., Nishino, S., and Itoh, M.: Penetration of the 1990s warm temperature anomaly of the Atlantic Water in the Canada Basin, *Geophys. Res. Lett.*, 31, L20301, doi:10.1029/2004GL020860, 2004.

15 Shimada, K., Itoh, M., Nishino, S., McLaughlin, F. A., Carmack, E., Proshutinsky, A.: Halocline structure in the Canada Basin of the Arctic Ocean, *Geophys. Res. Lett.*, 32, L03605, doi:10.1029/2004GL021358, 2005.

Smethie, W. M., Schlosser, P., and Bönisch, G.: Renewal and circulation of intermediate waters in the Canadian Basin observed on the SCICEX 96 cruise, *J. Geophys. Res.*, 105, 1105–1121, 2000.

20 Smith, J. N., Ellis, K. M., and Boyd, T.: Circulation features in the central Arctic Ocean revealed by nuclear fuel reprocessing tracers from Scientific Ice Expeditions 1995 and 1996, *J. Geophys. Res.*, 104, 29663–29667, 1999.

Stigebrandt, A.: A model for the thickness and salinity of the upper layers of the Arctic Ocean and the relation between the ice thickness and some external parameters, *J. Phys. Oceanogr.*, 11, 1407–1422, 1981.

25 Stigebrandt, A.: The North Pacific: a global-scale estuary, *J. Phys. Oceanogr.*, 14, 464–470, 1984.

Swift, J. H., Jones, E. P., Carmack, E. C., Hingston, M., Macdonald, R. W., McLaughlin, F. A., and Perkin, R. G.: Waters of the Makarov and Canada Basins, *Deep-Sea Res. Pt. II*, 44, 1503–1529, 1997.

30 Timmermanns, M.-L., Garrett, C., and Carmack, E.: The thermohaline structure and evolution of the deep waters in the Canada Basin, *Arctic Ocean, Deep-Sea Res. Pt. I*, 50, 1305–1321, doi:10.1016/S0967(03)00125-0, 2003.

Arctic Ocean
Circulation

B. Rudels

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean
Circulation

B. Rudels

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Timofeyev, V. T.: Water Masses of the Arctic Basin, Gidromet. Izdat., Leningrad, 1960.
- Walsh, D. and Carmack, E. C.: The nested structure of Arctic thermohaline intrusions, Ocean Model., 5, 267–289, 2003.
- Werenskiold, W.: Coastal Currents, Geophysical publications, 10(13), Oslo, 1935.
- Weyl, P. K.: The role of the oceans in climate change, Meteor. Mon. 8, 37–62, 1968.
- 5 Wüst, G.: Relief und Bodenwasser im Nordpolarbecken, Zeitschrift für Erdkunde zu Berlin, 5/6, Zeitschrift der Gesellschaft fr Erdkunde zu Berlin, 163–180, 1941.

Arctic Ocean
Circulation

B. Rudels

Table 1. Properties of the Atlantic water, the Polar surface water and the deep water for different percentages of entrained ambient (Atlantic) water.

Entrained water %	Polar surface water			Atlantic water			Outflowing deep water		
	Temp	Sal	Density	Temp	Sal	Density	Temp	Sal	Density
0	-1.9	34.27	27.56	3.1	35	27.85	-1.9	35	28.15
20	-1.9	34.27	27.56	3.1	35	27.85	-0.9	35	28.09
40	-1.9	34.27	27.56	3.1	35	27.85	0.1	35	28.03
60	-1.9	34.27	27.56	3.1	35	27.85	1.1	35	27.97
80	-1.9	34.27	27.56	3.1	35	27.85	2.1	35	27.91

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean
Circulation

B. Rudels

Table 2. Transports of Atlantic water, Polar surface water and deep water for different amount of Polar surface water transformed on the shelves and different percentages of entrained ambient (Atlantic) water.

Ent	Polar surface water transport				Atlantic water transport				Deep water outflow			
	Water to shelves				Water to shelves				Water to shelves			
%	0	0.24	0.48	0.72	0	0.24	0.48	0.72	0	0.24	0.48	0.72
0	0.96	0.72	0.48	0.24	0.94	0.94	0.94	0.94	0	0.24	0.48	0.72
20	0.96	0.72	0.48	0.24	0.94	1.03	1.06	1.1	0	0.30	0.60	0.90
40	0.96	0.72	0.48	0.24	0.94	1.10	1.26	1.44	0	0.40	0.80	1.20
60	0.96	0.72	0.48	0.24	0.94	1.30	1.66	2.04	0	0.60	1.20	1.80
80	0.96	0.72	0.48	0.24	0.94	1.90	2.86	3.84	0	1.20	2.40	3.60

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean Circulation

B. Rudels




Fig. 1. A map by Silas Bent, 1872, showing the warm ocean current and their assumed impact on the ice cover of the Arctic Ocean. From Hayes (2003).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean Circulation

B. Rudels





Fig. 2. Upper panel, map of the Arctic Ocean showing the drift of Fram and the track of Nansen and Johansen, map by Nansen, reproduced in Hayes (2003). Lower panel: temperature and salinity section based on the hydrographic observations made from Fram during the drift from the Laptev Sea to Fram Strait. From Nansen (1902).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

Back

Close

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 3. A temperature section from the Nordic Seas into the Arctic Ocean compiled by Wüst from available observations made between the 1880s and the 1930s. From Wüst (1941).

Arctic Ocean Circulation

B. Rudels




Fig. 4. Bathymetry of the Arctic Ocean (IBCAO) (Jakobsson et al., 2008), map from Rudels et al. (2011).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean Circulation

B. Rudels



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶|](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 5. Upper panel: schematics showing oceanic heat loss under freezing conditions and in the presence of sea ice (from Rudels, 1995). Lower panel: the evolution of the upper layer when warm water encounters sea ice under freezing conditions (from Rudels et al., 1999a).

Arctic Ocean Circulation

B. Rudels




Fig. 6. Potential temperature and salinity profiles and TS curves from different parts of the Arctic Ocean showing the effects of wind mixing and heat entrained and lost to ice melt and to the atmosphere at the entrance to the Arctic Ocean, of haline convection bringing cold water into and cooling the thermocline in the Nansen Basin, and of heat mixed from below into and stored in halocline in the interior of the Arctic Ocean.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean Circulation

B. Rudels




Fig. 7. Potential temperature and salinity profiles and TS diagrams showing the difference of the winter mixed layers and the Atlantic cores of the Fram Strait branch (red) and the Barents Sea branch (blue). In spite of having a cooler and less saline Atlantic layer the winter mixed layer of the Barents Sea branch, created by stirring melt water into the upper part of the Atlantic water, is more saline than that of the Fram Strait branch. The filled diamonds indicate the salinity of the mixed layer if all oceanic heat is going to ice melt, the open diamonds the salinity if minimum heat is going to ice melt based on the temperature and salinity of the Atlantic water.

Arctic Ocean Circulation

B. Rudels




Fig. 8. Halocline features in the Arctic Ocean. The grey, dark red and magenta arrows indicate the temperature minimum created by the winter homogenization. The red arrows show on one station, the upper temperature maximum due to seasonal heating, the temperature maximum due to the inflow of Bering Strait Summer Water (BSSW) and a temperature minimum due to the presence of an (isolated?) eddy of Fram Strait branch lower halocline water. The Green arrows show the temperature maximum of the BSSW and the temperature minimum of the Bering Sea Winter Water (BSWW). Note that the BSSW is present also at the red upper seasonal temperature maximum on the green station and the temperature minimum of the winter homogenization is present above the BSSW maximum.

Arctic Ocean
Circulation

B. Rudels




Fig. 9. Schematics showing the formation and circulation of the waters feeding the Fram Strait branch and the Barents Sea branch lower haloclines and how they become covered by less saline shelf water and Pacific water. Based on Rudels et al. (2004), figure from Rudels (2009).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean
Circulation

B. Rudels




Fig. 10. Potential temperature and salinity profiles and TS curves showing the characteristics of some water masses in the Arctic Mediterranean. Green station, Greenland Sea, red station, Fram Strait, orange station, Nansen Basin slope, magenta station, Gakkel Ridge, cyan station, Amundsen Basin, black station, Makarov Basin, blue station Canada Basin. AW (Atlantic Water) AAW (Arctic Atlantic Water), RAW (Recirculating Atlantic Water), uPDW (upper Polar Deep Water) AIW (Arctic Intermediate Water), CBDW (Canadian Basin Deep Water) EBDW (Eurasian Basin Deep Water) NDW (Nordic Seas Deep Water) (/) indicates unstable stratification in either temperature or salinity (/) indicates stable stratification in both components. From Rudels (2009).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Arctic Ocean
Circulation**

B. Rudels




Fig. 11. Circulation of the Atlantic and intermediate waters in the Arctic Mediterranean Sea showing the mixing area north of Severnaya Zemlya and the gradual changes of the properties of the Atlantic water. From Rudels et al. (2011).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean
Circulation

B. Rudels




Fig. 12. Water masses mixing and interleaving in the Nansen Basin north of the Eastern Kara Sea. Central panel: temperature and salinity sections from the shelf to the Gakkel Ridge showing the warm saline Fram Strait Atlantic core and the cooler and less saline Barents Sea branch water on the slope and also on the basin side of the warm core. Upper panel: potential temperature and salinity profiles and TS curves from the deep basin showing the interleaving structures and different fronts. Lower panel: potential temperature and salinity profiles and TS curves from the slope showing the characteristics of the two branches and the mixing between them across the front. Data from Polarstern 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean
Circulation

B. Rudels




Fig. 13. Water masses mixing and interleaving along the Gakkel Ridge north of the Laptev Sea. Central panel: temperature and salinity sections from the shelf along the Gakkel Ridge showing a colder and less saline Fram Strait Atlantic core compared to the Kara Sea section (Fig. 12) and the Barents Sea branch water has spread from the slope into the basin. It is also on the basin side of the warm core. Upper panel: potential temperature and salinity profiles and TS curves from the deep basin showing the interleaving structures. Lower panel: potential temperature and salinity profiles and TS curves from the slope showing interleaving between the two branches. Note that no smooth warm and saline Atlantic core is present. Interleaving layers are found everywhere. Data from Polarstern 2007.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean
Circulation

B. Rudels




Fig. 14. Potential temperature and salinity profiles and TS curves comparing the Barents Sea branch characteristics with those found in the other basins. The Barents Sea branch could, at least theoretically, supply the Atlantic and intermediate water to the Arctic Ocean Basins beyond the Nansen Basin and the Gakkel Ridge. Data from Polarstern 2007 and Oden 2005.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean Circulation

B. Rudels



[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I ▲](#)[I ▼](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fig. 15. Cold and brine enriched saline water observed west of Novaya Zemlya observed in 1901. From Knipovich (1905).

Arctic Ocean Circulation

B. Rudels




Fig. 16. Schematics showing to the left the evolution of the characteristics of dense shelf waters (A) due to entrainment as they sink down the continental slope (B). Depending upon the initial salinity the final characteristics form an envelope (C), which merges with the boundary current (E) to form the transformed water column (D). An idealised picture of plumes sinking to different depth is shown to the right. From Rudels and Quadfasel (1991).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean Circulation

B. Rudels




Fig. 17. Deep and bottom water characteristics from the Nansen Basin, Amundsen Basin, Makarov Basin and Canada Basin. Note the absence of a deep temperature minimum in the Makarov Basin and that the temperature minimum observed in the Canada Basin could derive from the Makarov Basin and the deep salinity maximum in the Amundsen Basin could be caused by Makarov Basin deep water crossing the Lomonosov Ridge.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Arctic Ocean
Circulation

B. Rudels




Fig. 18. Potential temperature and salinity profiles and TS curves from stations in the Makarov Basin, the Amundsen Basin and the intra-basin in the Lomonosov Ridge (lower map). In the intra-basin the red station in the interior and the green station at the sill, as well as the stations in the Amundsen and Makarov basins, are taken in 2005 by Oden. The magenta and yellow (not shown in the profiles and TS curves) are taken by Oden in 2001. The 2001 stations indicate that water deriving from the Amundsen Basin deep water was present in the deepest layer, located above the sill depth to the Makarov Basin. If this water crossed the sill it could sink down to 2500 m in the Makarov Basin.

Arctic Ocean
Circulation

B. Rudels





Fig. 19. Sections of dynamical heights across the Lomonosov Ridge with the Makarov Basin to the right and the Amundsen Basin to the left taken from Polarstern 1996 (upper panel) and Oden 2005 (lower panel). In 2005 the dynamic height at 2000 db was higher in the Amundsen Basin then in the Makarov Basin indicating a negative pressure gradient at sill depth directed from the Makarov Basin into the Amundsen Basin that could block the inflow of Amundsen Basin deep water into the Makarov Basin. In 1996 the situation was reversed.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean
Circulation

B. Rudels




Fig. 20. Potential temperature and salinity sections from Fram Strait taken in 1997 by R/V *Lance*. The West Spitsbergen Current and the East Greenland Current are indicated by red and blue ellipses, respectively.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 21. Idealised baroclinic exchanges through Fram Strait, where the upper layer in the East Greenland Current is created by allowing 0.02 Sv of ice melt on the Atlantic water. The salinity of the low salinity layer, 34.27, is determined from the constraint that minimum heat goes to ice melt. The Atlantic water has temperature 3.1 and salinity 35. Only one pressure reversal in created and nothing, except the rise in sea level, will stop the inflow of Atlantic water and generate a barotropic component to the outflow.

Arctic Ocean
Circulation

B. Rudels




Fig. 22. Idealised baroclinic exchanges through Fram Strait. Half of the low salinity upper layer is assumed advected back on the shelves, where ice formation and brine rejection increase the salinity to 35 and the shelf water sinks down the slope entraining ambient (Atlantic) water. The increased density in the lower layer causes a second pressure reversal to form and a deep outflow is generated. However, the deep outflow does not cease, when mass balance is reached and the sea level in the Arctic Ocean will drop, adding a barotropic component to the Atlantic inflow to conserve mass.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Arctic Ocean
Circulation

B. Rudels





Fig. 23. The idealised transports in the boundary current shown together with the observed barotropic eddies and the recirculation occurring in the central part of Fram Strait (from Schauer et al., 2008). Does such idealised exercise contribute to our understanding and how important are the exchanges in the central part of Fram Strait for the water mass transformations in the Arctic Ocean and the climate of the Arctic?

Title Page	Abstract	Introduction
Conclusions	References	
Tables	Figures	
◀	▶	
◀	▶	
Back	Close	
Full Screen / Esc		

[Printer-friendly Version](#)[Interactive Discussion](#)

**Arctic Ocean
Circulation**

B. Rudels

**Fridtjof Nansen****Bjørn Helland-Hansen****Johan W. Sandström****Martin Knudsen****V. Walfrid Ekman**

Discussion Paper | Discussion Paper

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)