

List of Alterations (relatively to the accepted manuscript)

Dear Sirs,

5 The final version of the manuscript has minor alterations relative to the previous revised version (accepted for publication),
namely:

Fig.2 – The colours of some of the data points in Fig. 2b have been corrected, however these changes did not imply any
change in the statistical analysis.

10

15

20

The following pages show the alterations, in the form of marked manuscript (alterations highlighted in yellow).

Technical Note: Algal Pigment Index 2 in the Atlantic off the Southwest Iberian Peninsula: standard and regional algorithms

5 Priscila Costa Goela^{1,2}, Sónia Cristina^{1,2}, Tamito Kakiyama³, John Icely^{1,2}, Gerald Moore⁴, Bruno Fragoso^{1,2}, Alice Newton^{1,5}

¹Centre for Marine and Environmental Research, FCT, University of Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

²Sagremarisco Lda., Apartado 21, 8650-999 Vila do Bispo, Portugal

³University of Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

10 ⁴Bio-Optika, Crofters, Gunnislake, UK

⁵Norwegian Institute for Air Research-IMPEC, Box 100, 2027 Kjeller, Norway

Correspondence to: Priscila Costa Goela (priscila.goela@gmail.com)

Abstract. In this study, Algal Pigment Index 2 (API2) is investigated in Sagres, an area located in the Atlantic off the southwestern Iberian Peninsula. Standard results provided by MEdium Resolution Image Spectrometer (MERIS) ocean
15 **colour** sensor were compared with alternative data products, determined through a regional inversion scheme, using both MERIS and in situ remote sensing reflectances (R_{rs}) as input data. The reference quantity for performance assessment is in situ total chlorophyll *a* (TChl*a*) concentration estimated through phytoplankton absorption coefficient (i.e., equivalent to API2). Additional comparison of data products has also been addressed **for** TChl*a* concentration determined by High Performance Liquid Chromatography. The MERIS matchup analysis revealed a systematic underestimation of TChl*a*, which
20 was confirmed with an independent comparison of product maps analysis. The study demonstrates the importance of regional algorithms for the study area that could complement upcoming standard results of the **current** Sentinel-3/OLCI space mission.

Keywords: absorption, MERIS Algal Pigment Index 2, Multi-Layer Perceptron (MLP) neural nets, ocean **colour**, remote sensing.

25 1 Introduction

The MERIS space sensor, operated by the European Space Agency (ESA) on-board of the Envisat platform from 2002 to 2012, has been continuously supported by investigations for the assessment and improvement of data products. Commissioned studies include the validation of radiometric data such as the R_{rs} (Cristina et al., 2014; Kajiya et al., 2014), as well the analyses of derived product maps (Kajiya et al., 2014; D'Alimonte et al., 2014; Cristina et al., 2016). These
30 MERIS validation activities have established an important basis to address Earth Observation (EO) capabilities through the

Ocean Land Colour Instrument (OLCI) sensor launched on the Sentinel-3 satellite in February 2016. OLCI data products are the main component of the Copernicus European programme to monitor the marine environment, and the retrieval of Chlorophyll *a* (Chl_a) is a core task of the Sentinel-3 space mission. Chl_a is needed to estimate the phytoplankton biomass in the ocean and to contribute to a variety of inter-related investigations and applications, including climate data records, environmental legislation, and a number of economic activities such as fisheries and aquaculture. After the removal of the atmospheric contribution to the signal recorded at the top of the atmosphere, Chl_a can be estimated from the bottom-of-atmosphere (BOA) R_{rs} values using the standard approach with polynomial algorithms based on band-ratios of the input radiometric quantities. The corresponding MERIS data product is denoted Algal Pigment Index 1 (API1) (Morel and Antoine, 2011). The use of band-ratio is based on the assumption that seawater optical properties are driven by Chl_a. A tendency to overestimation has however been documented in optically complex marine conditions (D'Alimonte et al., 2014). This can occur when optically active constituents such as Coloured Dissolved Organic Matter (CDOM) and detrital particulate matter exceed their typical levels. The Chl_a retrieval accuracy declines in these optically complex conditions because the band-ratio approach attributes variations of the R_{rs} spectral slope to changes of Chl_a. In such cases, regionalized bio-optical algorithms are required (Bricaud et al., 2002, Gregg and Casey, 2004). Alternative ocean colour inversion schemes adopted to improve the Chl_a retrieval from space include artificial Neural Nets (NN) using R_{rs} at selected wavelengths as input. In the case of MERIS standard deliverables, this corresponds to the API2 data product (Doerffer and Schiller, 2007).

Although NNs can in principle model any relationship between apparent and inherent optical properties, their performance is in practice mostly determined by the dataset used for their training. Specific analyses are then needed to compare the standard MERIS API2 results with independent estimates. This main requirement is addressed in the present work by: 1) developing and assessing the performance of an independent regional Multilayer Perceptron (MLP) scheme to retrieve results equivalent to MERIS API2 values; and 2) comparing MERIS standard and regional API2 product maps.

The region under study is the Atlantic off the Southwestern Iberian Peninsula, where in situ reference data were collected at three stations off the Sagres region at 2, 10 and 18 km from the coast (henceforth stations A, B and C, respectively). The study is conducted based on both matchup analyses and product map inter-comparisons, with timely presentation of the results acknowledging, not only the planned MERIS data reprocessing, but also the need for a benchmark for the analysis of the upcoming OLCI API2 deliverables. An added value of this study is to confirm that qualitative evaluations based on product maps comparison can complement matchup data at the early mission stages of OLCI, when the statistical significance of matchup analysis is limited.

30 **2 Data and Methods**

Field campaigns were performed from 2008 to 2012 at the three study sites, with simultaneous collection of water samples and radiometric measurements. MERIS Level 2 Full Resolution (FR, 290 m x 260 m) and Reduced Resolution (RR, 1.20 km

x 1.04 km) satellite images were extracted for matchup analysis and product map comparison, respectively, and analyzed with the Basic ERS & ENVISAT (A) ATSR and MERIS Toolbox (BEAM version 4.9). The MEGS 8.1 processor (MERIS third reprocessing) was used to derive level 2 data, in agreement with previously reported extraction procedures (Cristina et al., 2014, 2015). The selection of satellite images was restricted to images without clouds and contamination, as indicated by not having specific Product Confidence (PCD), sun glint and ice flags. More details on the image selection criteria and full description of flags are reported in Cristina et al. 2016. TChl a concentration (monovinyl Chl a + divinyl Chl a + chlorophyllide a + phaeopigments) was determined by High Performance Liquid Chromatography (HPLC), according to Wright and Jeffrey (1997), herein referred to as TChl $a_{\text{HPLC}}^{\text{REF}}$. The protocols adopted for TChl a extraction, identification and quantification procedures are reported in Goela et al. (2014, 2015).

10 2.1 In situ reference data

In situ radiometric measurements were acquired with a tethered attenuation coefficient chain sensor (TACCS, Satlantic[®]), supporting a hyperspectral surface irradiance sensor $E_s(\lambda)$ and a subsurface radiance sensor $L_u(\lambda)$, as well as a tethered attenuation chain equipped with four irradiance sensors at nominal depths of 2, 4, 8 and 16 m. Normalized water leaving reflectance (ρ_N) was computed with Eq. (1):

$$15 \quad \rho_N(\lambda) = \pi \frac{L_w(\lambda)}{E_s(\lambda)}, \quad (1)$$

where L_w is the water leaving radiance determined by propagating L_u from below to above the sea surface and corrected for self-shading following (Gordon and King, 1992). $\rho_N(\lambda)$ corresponds to the remote sensing reflectance R_{rs} upon scaling with π .

For the determination of in situ absorption of phytoplankton pigments at 442 nm ($a_{\text{ph}}(442)$), seawater filtrates (0.5 L) were collected on GF/F filters (pore size 0.7 μm), which were then analyzed with the transmittance-reflectance technique as in (Tassan and Ferrari, 2002), using a dual beam-spectrophotometer (GBC[®] CINTRA 40), equipped with an integrating sphere. The phytoplankton absorption was determined as the difference between the total particulate and detrital absorption, which were measured before and after sodium hypochlorite bleaching (Ferrari and Tassan, 1999; Goela et al., 2013), respectively. The API2 in situ equivalent algal pigment index TChl $a_{\text{ABS}}^{\text{REF}}$ was then estimated by converting $a_{\text{ph}}(442)$ into API2, using the same regression coefficients presented in Sect. 2.2.2.

2.2 Chlorophyll a retrieval algorithms

2.2.1 MERIS Standard algorithm API2

This standard product is estimated with two NNs. The first NN computes BOA R_{rs} values by removing the atmospheric radiometric contribution from input space-born R_{rs} values. The second NN utilizes the BOA R_{rs} to derive the $a_{\text{ph}}(442)$. The

final API2 product is then computed as $MER^{API2} = A \times a_{ph}(442)^B$, with power-law regression coefficients $A=21.0$ and $B=1.04$ derived from field measurements in the German Bight and Norwegian waters (Doerffer and Schiller, 2007).

2.2.2 Regional MLP NN algorithm

The regional MLP for retrieving the data product equivalent to API2 has been trained with the in situ data collected at the
 5 Sagres site (instructions for independent implementation by users are provided as supplemental material
<http://ocportugal.org/sites/default/files/mlpSgrAPI2.pdf>). This MLP is here applied to two different sets of input data for
 assessment of performance and for comparison of results. The first set consists of the in situ R_{rs} values (R_{rs}^{SITU}), and the
 second set includes standard MERIS BOA R_{rs} data (R_{rs}^{MER}). Corresponding data products are denoted $MLP(R_{rs}^{SITU})$ and
 $MLP(R_{rs}^{MER})$, respectively. In both cases, R_{rs} at 490, 510 and 560 nm were selected as input channels, in agreement with the
 10 reference study (Cristina et al., 2014).

A novelty detection scheme (D'Alimonte et al., 2014; Bishop, 1994) was used to verify the algorithm applicability range by
 evaluating the representativeness of the input data in the training dataset (D'Alimonte et al., 2003; Mélin et al., 2011; Sá et
 al., 2015). The adopted applicability range is based on a novelty index (η) presented in published works (D'Alimonte et al.,
 2013; Sá et al., 2015). A revision is however applied for the scope of this work. This updated version considers all
 15 dimensions of the Principal Component Analysis (PCA) of selected input data, rather than only the first three components
 considered in the past (see the supplemental material for details). This updated definition is more effective for cases where
 the variability of training and application data tends to occur at different wavelengths (details not presented here). Key
 features are: 1) η is bounded between 0 and ∞ ; 2) the more the R_{rs} spectrum is similar to the in situ MLP training
 measurements, the lower is its η ; and 3) an R_{rs} spectrum is considered within the MLP applicability range when $\eta \leq 1$.

20

3 Results

The main tasks of this study are: 1) to evaluate the performance of regional MLP algorithm and the MER^{API2} results with
 respect to the in situ $TChla_{ABS}^{REF}$ reference measurements; 2) to verify the applicability of the regional $MLP(R_{rs}^{MER})$ and to
 compare product maps with MER algal pigment indices; and 3) to extend the analysis by also considering $TChla_{HPLC}^{REF}$ for
 25 data product assessment.

The statistical figures used to evaluate the estimated (y) in relation to the reference in situ $TChla(x)$, were absolute (ϵ) and
 signed (δ) percent differences, defined as:

$$\epsilon = \frac{1}{N} \sum_{i=1}^N \frac{|y_i - x_i|}{x_i} \times 100; \quad \delta = \frac{1}{N} \sum_{i=1}^N \frac{y_i - x_i}{x_i} \times 100, \quad (2)$$

where N is the total number of samples and i is the sample index. For product maps comparison, the absolute (ϵ^*) and signed
 30 (δ^*) unbiased differences were instead determined as:

$$\varepsilon^* = \frac{1}{N} \sum_{i=1}^N \frac{|y_i - x_i|}{y_i + x_i} \times 200; \delta^* = \frac{1}{N} \sum_{i=1}^N \frac{y_i - x_i}{y_i + x_i} \times 200, \quad (3)$$

where x_i and y_i are the $MLP(R_{rs}^{MER})$ and MER^{API2} values, respectively, taking the mean of the two values as a reference. In addition, the coefficient of determination r^2 between the evaluated quantities is also reported. The total number of samples used to validate MER^{API2} and $MLP(R_{rs}^{MER})$ algorithms results with respect to the in situ reference measurements was $N=54$.

5 In contrast, the total number of samples for assessing the performance of regional MLP algorithm with in situ reference measurements ($MLP(R_{rs}^{SITU})$), was $N=297$. This larger number of samples is based on the data from 4-8 radiometric casts for each in situ TChla sample at each location.

3.1 Matchup data analysis

The top panels of Fig. 1 present the matchup comparisons of MER^{API2} , $MLP(R_{rs}^{MER})$ and $MLP(R_{rs}^{SITU})$ with respect to the in situ reference $TChla_{ABS}^{REF}$ (Figs. 1a, 1b and 1c, respectively). While MER^{API2} underestimated TChla ($\delta = -34\%$) especially at higher concentrations, the regional products slightly overestimated TChla ($\delta = 11\%$ for $MLP(R_{rs}^{MER})$ and 2% for $MLP(R_{rs}^{SITU})$). The best agreement between data sets was obtained with $MLP(R_{rs}^{SITU})$, while MER^{API2} showed larger uncertainties. Table 1 presents the matchup analysis where the underestimation of MER^{API2} in relation to TChla is relatively constant (35%, 32% and 34%, in stations A, B and C, respectively) in all stations, but the correlation coefficient improves with distance offshore (0.22, 0.60, 0.67 in stations A, B and C, respectively).

In general, the matchup analysis with $TChla_{HPLC}^{REF}$ revealed higher uncertainties for MER^{API2} , $MLP(R_{rs}^{MER})$ and $MLP(R_{rs}^{SITU})$, as detailed in Fig. 1 (lower panel). Note that also in this case $MLP(R_{rs}^{SITU})$ presented the best results, with the highest coefficient of determination and the lowest bias. Similar to what was documented for $TChla_{ABS}^{REF}$, the bias for $TChla_{HPLC}^{REF}$ displayed only small differences between the sampling stations. The coefficient of determination instead increased from station A to station C. The underestimation of MER^{API2} in relation to $TChla_{HPLC}^{REF}$ was also observed, but with a lower bias (Fig. 1d). These observations are schematized in Fig. 2, where MER^{API2} was considered as the baseline. A complementary comparison with MER^{API1} is also presented for completeness. Results indicated an overestimation by the API1 algorithm in relation to both estimations of TChla (details not shown). The tendency of $TChla_{ABS}^{REF}$ to produce higher values than $TChla_{HPLC}^{REF}$ was also confirmed.

25 3.2 Comparison of product maps

The comparison of MERIS API2 standard product with the MLP regional results is presented on Fig. 3. The maps for the regional MLP (Fig. 3a) and the MER^{API2} (Fig. 3b) are shown in the top panel, together with the difference between MER^{API2} and $MLP(R_{rs}^{MER})$ shown in Fig. 3c. Overestimations of more than 35% in relation to the regional MLP are coloured in pink, and underestimations below 35 % are coloured in yellow, while differences between -35% and 35% are in green. The

MLP(R_{rs}^{MER}) region of applicability is shown in Fig. 3d, with black contours indicating the threshold $\eta=1$. Results indicate an underestimation by MER^{API2} of more than 35% in a significant part of the applicability range, especially near the coast.

The results from the application of Sagres regional MLP to the Atlantic off the Portuguese coast is presented in Fig. 3e and Fig. 3f. Besides the Sagres area (#3, in blue), two other regions of interest (ROIs) were chosen for comparison of product maps: Figueira da Foz (#1, in red) and Lisbon region (#2, in green, Fig. 3e). Note that ROI #1 and #2 have been selected for their contrasting features: the first influenced by Mondego river plume and the second by the Tagus estuary. The comparison between the MER^{API2} and regional MLP products is presented as a scatter plot (Fig. 3f), following the same colour coding of the three ROIs. The underestimation tendency of MER^{API2} in relation to in situ TChla was confirmed through this analysis. The results also indicated more pronounced differences in Mondego and Tagus ROIs, where values of TChla were higher.

The statistical figures of the product map comparison between MER^{API2} and regional MLP are summarized in Table 2. The applicability of the Sagres MLP was verified with the novelty detection scheme. The number of total and valid (i.e., $\eta < 1$) data points are denoted as N_{tot} and N_{val} , respectively. The Sagres ROI presents the highest number of valid data points, while Tagus region had the highest percentage of novel data points.

4 Discussion

This study analyzed the standard MERIS API2 product by considering the TChla retrieval in the coastal waters of Portugal. Data product comparisons have been performed by developing and applying a regional MLP trained with Sagres in situ data and accounting for its applicability range. The work highlighted a tendency of MER^{API2} to underestimate TChla, not only when the reference values were derived through $a_{ph}(442)$ but also when determined by HPLC. This result is consistent with other studies addressing low productivity waters (Tilstone et al., 2012). This underestimation tendency is more pronounced at higher concentrations but not observed in the results of the regional MLP. Possible explanations can be uncertainties in BOA R_{rs} values, as well as in specific properties of the NN inversion scheme used to compute the standard API2 values. It is noted that the MERIS NN scheme for API2 retrieval is scoped for global applications in both Case 1 and optically complex waters. This general applicability might limit the algorithm performance in the presence of specific bio-optical relationships at the regional scale. An example could be the upwelling along the coast of Portugal (Loureiro et al., 2005; Goela et al., 2015).

As a contribution to the forthcoming OLCI mission, the present work also provides indications to enhance standard OLCI API2 results by including additional training samples in the synthetic dataset used for the development of the MERIS NN scheme. The overestimation of $TChla_{ABS}^{REF}$ in relation to $TChla_{HPLC}^{REF}$ has been identified in this study as one of the reasons for the systematic difference observed in the comparison of MER^{API2} with both in situ referred targets (Fig. 2b).

The regional MLP using in situ R_{rs} as input produced highly accurate results (bias of 2%), when relating R_{rs}^{SITU} to reference measurements of $TChla_{ABS}^{REF}$. When MERIS R_{rs} is used, the bias is slightly higher, probably due to the uncertainties of the atmospheric correction (Cristina et al., 2014). It is also reported that a cross-validation analysis performed by splitting the in

situ data in different subsets to develop and assess the regional MLP documented an increase from 2 to 9% of the bias (details not presented). As observed for the standard NN inversion schemes, the performance of the regional MLP could be enhanced through a better representation of the optical properties of the study region: the collection of additional field measurements is hence recommended. Another aspect that has been considered is the reduction in bias when the training dataset was $TChla_{ABS}^{REF}$ estimated with a_{ph} at 440 nm (7% of bias). This indicates that the specific selection of the wavelength of the maximum phytoplankton absorption could allow for a better TChla parameterization and hence also lead to a more accurate regional MLP.

The strong relationship between R_{rs} and the phytoplankton coefficient of absorbance at 442 nm suggests the presence of Case 1 waters. The better agreement with $TChla_{ABS}^{REF}$ rather than with $TChla_{HPLC}^{REF}$ can be explained by considering that the training of the neural net was performed with $TChla_{ABS}^{REF}$. An additional explanation could be that $TChla_{ABS}^{REF}$ was determined using $a_{ph}(442)$, which is likely better related to R_{rs} than $TChla_{HPLC}^{REF}$ (both $a_{ph}(442)$ and R_{rs} directly represent optical properties). A caveat would however apply to this argument, whereas $TChla_{HPLC}^{REF}$ is a direct measurement of the TChla concentration, $TChla_{ABS}^{REF}$ is an indirect measurement which has errors associated with the laboratory determination of $a_{ph}(442)$.

It is also noted that the regional relationship between a_{ph} at 442 nm and TChla retrieved by HPLC is close to that used in MER^{API2} ($TChla_{MERIS} = 21 a_{ph}(442)^{1.04}$, $TChla_{SAGRES} = 27 a_{ph}(442)^{1.13}$). However, the local relationship between TChla and $a_{ph}(442)$ corresponds to a coefficient of determination $r^2 = 0.8$. Hence, about 20% of variability of TChla is not related to $a_{ph}(442)$.

The ROIs data analysis indicates lower MERIS API2 values with respect to equivalent results derived with the regional MLP, especially when the TChla concentration increases. This finding is in a good agreement with the matchup results, thereby, highlighting the benefit of independent comparison of product maps to qualitatively evaluate data products at an early stage of ocean colour space missions (e.g., OLCI).

5 Conclusions

The scope of this technical note was to analyze the MERIS standard API2 product in the Southwestern coast of Portugal. A regional MLP algorithm to retrieve TChla, estimated through phytoplankton absorption coefficient, was implemented and applied for this purpose. This regional algorithm produced good agreement with in situ data, hence indicating a high accuracy of regional MLP products. The applicability of the regional MLP in the study area was verified by a novelty detection scheme. With this information, the study reports an underestimation tendency of MER^{API2}, which is consistent with other European basins within low ranges of this constituent. The results of the regional MLP were closer to the in situ reference for API2 – TChla estimated with $a_{ph}(442)$ – than to TChla determined by HPLC. This work also indicates that the use of a regional relationship between phytoplankton absorption and pigment concentration is expected to improve the accuracy of global ocean colour remote sensing products.

This study has highlighted the usefulness of maintaining in situ measurement programmes for validation purposes of ongoing ocean colour missions. Moreover, it has also demonstrated the importance of developing regional algorithms that not only complement standard approaches, but that can also be applied for the qualitative data assessments of new ocean colour missions in the early stages of product map delivery (e.g., Sentinel-3).

5 Data availability

The majority of the in situ data used in this work can be accessed through the ESA MERIS MAtchup In-situ Database (<http://mermaid.acri.fr/home/home.php>) and the MERIS satellite data can be accessed through the Optical Data processor of ESA (http://www.odesa-info.eu/process_basic/basic.php).

Acknowledgements

- 10 The authors thank Dr. Davide D'Alimonte for his contribution to the MLP NN algorithm development and training, and wise advice both on the methodology design and in the interpretation of the results. This work was supported in part by the European Space Agency (ESA) for the “Technical Assistance for the Validation of MERIS Marine Products at Portuguese oceanic and coastal sites”(contract no. 21464/08/I-O) and “MERIS validation and algorithm 4th reprocessing” (contract no. ARG/003-025/14067Sagremarisco and ARG/003-025-1406/CIMA). Priscila Costa Goela and Sónia Cristina were funded by
15 PhD grants from the Portuguese FCT(SFRH/BD/78356/2011 and SFRH/BD/78354/2011, respectively); Alice Newton was funded by EU FP7 project DEVOTES (grant no. 308392); John Icely is funded by EU FP7 AQUA-USER (grant no. 607325), and Horizon 2020 AquaSpace (grant no. 633476).

References

- 20 Bishop, C. M.: Novelty Detection and Neural Network Validation, *IEE Proc. Vision and Image & Sig. Proc.*, 141, 217–222, doi:10.1049/ip-vis:19941330, 1994.
- Bricaud, A., Bosc, E and Antoine, D.: Algal biomass and sea surface temperature in the Mediterranean basin. Intercomparison of data from various satellite sensors, and implications for primary production estimates. *Remote Sensing of Environment*, 81, 163 – 178, 2002.
- Cristina, S., Cordeiro, C., Lavender, S., Goela, P.G., Icely, J., Moore, G. and Newton, A.: Seasonal-Trend decomposition time series based on Loess applied to MERIS products from the SW Iberian Peninsula: Sagres. *Remote Sensing*, 8(6), 449; doi:10.3390/rs8060449, 2016.
- 25 Cristina, S., D'Alimonte, D., Goela, P.G., Kakiyama, T., Icely, J., Moore, G., Fragoso, B.D.D. and Newton, A.: Standard and Regional Bio-optical Algorithms for Chlorophyll a Estimates in the Atlantic off the Southwestern Iberian Peninsula. *IEEE Geosci. Remote S.*, 13, 6, 757-761, DOI:10.1109/LGRS.2016.2529182, 2016.
- Cristina, S., Icely, J., Goela, P., DelValls, T. and Newton, A.: Using remote sensing as a support to the implementation of the European Marine Strategy Framework Directive in SW Portugal, *Cont. Shelf Res.*, 108, 169–177, doi:10.1016/j.csr.2015.03.011, 2015.
- 30 Cristina, S., Moore, G., Goela, P. C., Icely, J. and Newton, A.: In situ validation of MERIS marine reflectance off the southwest Iberian Peninsula: assessment of vicarious adjustment and corrections for near-land adjacency, *Int. J. Remote Sens.*, 35, 6, 2347–2377, doi:10.1080/01431161.2014.894657, 2014.
- D'Alimonte, D., Mélin, F., Zibordi, G. and Berthon, J.-F.: Use of the Novelty Detection Technique to Identify the Range of Applicability of Empirical Ocean Colour Algorithms, *IEEE Trans. Geosc. Rem. Sens.*, 41, 2833–2843, doi:10.1109/TGRS.2003.818020, 2003.
- 35 D'Alimonte, D., Zibordi, G., Kakiyama, T. and Berthon, J.-F.: Comparison between MERIS and regional high-level products in European seas, *Remote Sens. Environ.*, 140, 378–395, doi:10.1016/j.rse.2013.07.029, 2014.
- Doerffer, R. and Schiller, H.: The MERIS Case 2 water algorithm, *Int. J. Remote Sens.*, 28, 3-4, 517–535, doi:10.1080/01431160600821127, 2007.

- Ferrari, G. and Tassan, S.: A method using chemical oxidation to remove light absorption by phytoplankton pigments, *J. Phycol.*, 35, 1090–1098, doi:10.1046/j.1529-8817.1999.3551090.x, 1999.
- Goela, P. C., Icely, J., Cristina, S., Danchenko, S., DelValls, T. A. and Newton, A.: Using bio-optical parameters as a tool for detecting changes in the phytoplankton community (SW Portugal), *Estuar. Coast. Shelf S.*, 167, part A, 125–137, doi:10.1016/j.ecss.2015.07.037, 2015.
- 5 Goela, P. C., Icely, J., Cristina, S., Newton, A., Moore, G. and Cordeiro, C.: Specific absorption coefficient of phytoplankton off the southwest coast of the Iberian Peninsula: A contribution to algorithm development for ocean colour remote sensing, *Cont. Shelf Res.*, 52, 119 – 132, doi:10.1016/j.csr.2012.11.009, 2013.
- Goela, P., Danchenko, S., Icely, J., Lubian, L., Cristina, S. and Newton, A.: Using CHEMTAX to evaluate seasonal and interannual dynamics of the phytoplankton community off the South-west coast of Portugal, *Estuar., Coast. Shelf S.*, 151, 112 – 123, doi:10.1016/j.ecss.2014.10.001, 2014.
- 10 Gordon, H. R. and Ding, K.: Self-shading of in-water optical instruments, *Limnol. Oceanogr.*, 37, 491–500, doi:10.4319/lo.1992.37.3.0491, 1992.
- Gregg, W.W. and Casey, H.W. Global and regional evaluation of the SeaWiFS chlorophyll data set. *Remote Sensing of Environment*, 93, pp. 463–479, doi:10.1016/j.rse.2003.12.012, 2004.
- Kajiyama, T., D’Alimonte, D. and Zibordi, G.: Match-up analysis of MERIS radiometric data in the Northern Adriatic Sea, *IEEE Geosci. Remote Sens. Lett.*, 11, 1, 19–23, doi:10.1109/LGRS.2013.2244844, 2014.
- 15 Loureiro, S., Newton, A. and Icely, J.: Microplankton composition, production and upwelling dynamics in Sagres (SW Portugal) during the summer of 2001, *Sci. Mar.*, 69, 3, 323–341, doi:10.3989/scimar.2005.69n3323, 2005.
- Mélin, F., Vantrepotte, V., Clerici, M., D’Alimonte, D., Zibordi, G., Berthon, J. F. and Canuti, E.: Multi-sensor satellite time series of optical properties and chlorophyll-*a* concentration in the Adriatic Sea, *Prog. Oceanogr.*, 91, 3, 229–244, doi:10.1016/j.pocean.2010.12.001, 2011.
- Morel, A. and Antoine, D.: Pigment index retrieval in case 1 waters, ATBD 2.9, ESA Doc. No. PO-TN-MEL-GS-0005 (version 4.3), 2011.
- 20 Sá, C., D’Alimonte, D., Brito, A., Kajiyama, T., Mendes, C., Vitorino, J., Oliveira, P., Silva, J. and Brotas, V.: Validation of standard and alternative satellite ocean-color chlorophyll products off Western Iberia, *Remote Sens. Environ.*, 168, 403 – 419, doi:10.1016/j.rse.2015.07.018, 2015.
- Tassan S. and Ferrari, G.: A sensitivity analysis of the “Transmittance-Reflectance” method for measuring light absorption by aquatic particles, *J. Plankton Res.*, 24, 8, 757–774, doi:10.1093/plankt/24.8.757, 2002.
- 25 Tilstone, G. H., Peters, S. W., Van Der Woerd, H. J., Eleveld, M. A., Ruddick, K., Schönfeld, W., Krasemann, H., Martinez-Vicente, V., Blondeau-Patissier, D., Röttgers, R., Sørensen, K., Jørgensen, P. V. and Shutler, J. D.: Variability in specific-absorption properties and their use in a semi-analytical ocean colour algorithm for MERIS in North Sea and Western English Channel Coastal Waters, *Remote Sens. Environ.*, 118, 320 – 338, doi:10.1016/j.rse.2011.11.019, 2012.
- Wright, S. and Jeffrey, S.: High-resolution HPLC system for chlorophylls and carotenoids of marine phytoplankton, *in* *Phytoplankton pigments in oceanography*. UNESCO Publ., 1997, 327–341.

30

Table 1. Comparison between standard (MER^{API2}), regional bio-optical algorithms ($MLP(R_{rs}^{MER})$ and $MLP(R_{rs}^{SITU})$) and $TChla^{REF}$

	N				$\varepsilon(\%)$				$\delta(\%)$				r^2			
	A	B	C	All	A	B	C	All	A	B	C	All	A	B	C	All
MER^{API2} vs $TChla_{ABS}^{REF}$	18	17	19	54	45	35	38	39	-35	-32	-34	-34	0.22	0.60	0.67	0.49
MER^{API2} vs $TChla_{HPLC}^{REF}$	18	17	19	54	48	39	42	43	-21	-24	-26	-24	0.18	0.54	0.66	0.38
$MLP(R_{rs}^{MER})$ vs $TChla_{ABS}^{REF}$	18	17	19	54	23	32	30	29	8	8	16	11	0.69	0.51	0.85	0.67
$MLP(R_{rs}^{MER})$ vs $TChla_{HPLC}^{REF}$	18	17	19	54	66	45	49	54	39	16	30	29	0.38	0.49	0.49	0.43
$MLP(R_{rs}^{SITU})$ vs $TChla_{ABS}^{REF}$	93	91	113	297	16	17	19	17	3	-4	7	2	0.88	0.91	0.91	0.91
$MLP(R_{rs}^{SITU})$ vs $TChla_{HPLC}^{REF}$	93	91	113	297	56	35	39	43	27	7	20	18	0.48	0.86	0.61	0.63

Table 2. Comparison between the regional $MLP(R_{rs}^{MER})$ and the standard MER^{API2} (The location of ROIs is presented in Fig. 3e).

ROI	N_{tot}	N_{val}	$\varepsilon^*(\%)$	$\delta^*(\%)$	r^2
#1	2122	2075	43	-43	0.70
#2	3383	1739	32	-30	0.71
#3	2946	2224	20	-15	0.76
Total	8451	6038	32	-29	0.76

Table 3. List of notations.

API1	Algal Pigment Index 1
API2	Algal Pigment Index 2
BEAM	Basic ERS & ENVISAT (A) ATSR and MERIS Toolbox
BOA	Bottom-of-atmosphere
CDOM	Coloured Dissolved Organic Matter
Chl a	Chlorophyll a
EO	Earth Observation
$E_s(\lambda)$	Surface downwelling incident irradiance
HPLC	High Performance Liquid Chromatography
$L_u(\lambda)$	Subsurface upwelling radiance
$L_w(\lambda)$	Water leaving radiance
MER ^{API2}	MERIS Algal Pigment Index 2 standard product
MERIS	MEdium Resolution Image Spectrometer
MLP	Multilayer Perceptron
MLP(R_{rs}^{MER})	Regional TChl a products computed using inversion schemes based on the MLP NN using standard MERIS BOA R_{rs}
MLP(R_{rs}^{SITU})	Regional TChl a products computed using inversion schemes based on the MLP NN using in situ R_{rs}
NN	Neural Nets
N_{tot}	Number of total (i.e., $\eta < 1$) data points
N_{val}	Number of valid (i.e., $\eta < 1$) data points
OLCI	Ocean Land Colour Instrument
PCA	Principal Component Analysis
r^2	Coefficient of determination
ROIs	Regions of interest
R_{rs}	Remote sensing reflectances
R_{rs}^{MER}	Standard MERIS BOA R_{rs}
R_{rs}^{SITU}	In situ R_{rs}
TChl a	Total Chlorophyll a
TChl a_{ABS}^{REF}	API2 in situ equivalent algal pigment index
TChl a_{HPLC}^{REF}	TChl a concentration (monovinyl Chl a + divinyl Chl a + chlorophyllide a + phaeopigments) determined by HPLC
δ	Signed percent differences
δ^*	Signed unbiased differences
ε	Absolute percent differences
ε^*	Absolute unbiased differences
η	Novelty index
ρ_N	Normalized water leaving reflectance




Figure 1: Comparison between MERIS standard Algal Pigment Index 2 and results obtained by applying the Multilayer Perceptron (MLP) regional **scheme** for the Sagres region. The top row panels present the matchup comparisons with respect to the in situ reference TChl a_{ABS}^{REF} , while the lower panels detail the matchup comparisons with TChl a_{HPLC}^{REF} .






Figure 2: a) Schematic diagram showing, respectively, underestimation and overestimation of MERIS algal pigment indices 1 and 2, relative to TChla, estimated through the absorption coefficient at 442 nm ($TChla_{ABS}^{REF}$) and measured by HPLC ($TChla_{HPLC}^{REF}$), and b) scatter plot of the $TChla_{ABS}^{REF}$ versus $TChla_{HPLC}^{REF}$.



5 **Figure 3: Comparison between Sagres regional Multilayer Perceptron (MLP) algorithm map with MERIS pigment index product map Algal Pigment Index 2 for the 25th August 2010, showing a) the product map of the regional MLP, b) standard API2 MERIS product map, c) difference between MER^{API2} and MLP(R_{rs}^{MER}), d) region of applicability of MLP(R_{rs}^{MER}), e) results of the application of the regional MLP to the Portuguese coast in the three regions of interest (shown in e). Please see Sect. 3.2 for a more detailed description of the panels. (Source: MER_RR_2PRAC20100825_103551_000026292092_00223_44365_0000.N1)**