

## Reply to Anonymous Referee #1

COMMENTS FROM REFEREE: I am happy with the modifications that give me more confidence that we are actually seeing quasi-stationary hydraulic controls. However, it could be taken one step further by just considering mass conservation. In a steady system with fully mixed downstream water, the downstream salinity must be determined by the salinities and relative volume fluxes of the two upstream layers (e.g. Arneborg, 2016, JGR, 121, 2035–2040). That could be mentioned at some parts of the text, as suggested in the detailed comments below. I also think that the discussion about Chl<sub>a</sub> and primary production could be extended somewhat, as also specified in more detail below.

Detailed comments:

Line 96-97: I still don't understand where this comes from. Prastewo et al (2009) find a mixing efficiency of about 0.11 which is less than the standard value of 0.2 used by most other researchers. Mixing may be small in sub-critical flow, but there is no reason to believe that the mixing efficiency is 10 times smaller than 0.11. In flows dominated by bottom friction, the bulk mixing efficiency may be smaller since most of the turbulent dissipation happens in the well mixed bottom boundary layer, but that is not really what you write.

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: We are having difficulties understanding what the referee finds to be a problem here. Our point is that mixing is much more intense in the supercritical part of the flow compared to the upstream, subcritical part. Existing studies have found the difference in mixing efficiency to be an order of magnitude, i.e.  $O(0.1)$  for supercritical flows and  $O(0.01)$  for subcritical flows. Is the referee concerned with the exact values of the efficiencies? We are deliberately not going into details with these because they are not so important. Therefore we are unable to see how our text is unclear about the point that we are trying to make. However, given the referee's criticism we are going to try to reformulate the two sentences in question.

COMMENTS FROM REFEREE: Line 186: Is it reasonable to assume that 80% of the volume flux takes place in the upper layer, as these results indicate? I think it is, but I also think this extra constraint on the results should be mentioned and evaluated, even though the evaluation cannot be very precise without velocity measurements.

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: This is a very good point that deserves attention. We will add both a general comment earlier in the manuscript, including a reference to Arneborg (2016), and a specific comment in the following paragraph.

COMMENTS FROM REFEREE: Lines 198-201: To me this is not that clear to see directly from Figs 1 and 2, but it may be an interpretation of the results. Please write some more explanation.

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: We will add some details to clarify this.

COMMENTS FROM REFEREE: Lines 234-236: Same as above. Is it reasonable to assume that 60% of the volume flux takes place in the upper layer?

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: This is a good point and question. We will add some text to answer it.

COMMENTS FROM REFEREE: Line 251: I had some problems understanding this. Maybe add “of the strait” to make clear what “upper part” refers to.

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: We agree and will modify the text.

COMMENTS FROM REFEREE: Line 255: Change to “...top of the slope.”

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: A typo. We will fix it.

COMMENTS FROM REFEREE: Line 256: Change to "...in case of a relatively..."

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: Another typo. which we will fix.

COMMENTS FROM REFEREE: Line 258: Change to "In nature, hydraulic controls..."

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: Yes, a comma is needed, which we will add.

COMMENTS FROM REFEREE: Lines 359-362: I understand this explanation for the upper layer in Fig. 8, where bottom water is mixed up into the upper layer. However, I would like some more text on how the authors interpret the enhanced Chl\_a in the lower layers. Why does entrainment of upper layer water enhance primary production in the lower layer? The nutrients are already there and the light is probably not better than upstream. Is it because primary producing plankton are mixed down into the layer where they can grow on the limited light sources there?

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: Yes, this is not clear. We will clarify the text.

COMMENTS FROM REFEREE: Lines 404-408: In my way of seeing this, it is the control that determines the downstream salinity through the relative volume fluxes in the two layers. The entrainment can do whatever it wants as long as it is strong enough to homogenize the fluid.

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: Yes, this is true. But our concern here is how the flow rates, and subsequently the water masses on the downstream side, are determined by the control. We see no reason to modify this part of the discussion.

COMMENTS FROM REFEREE: Line 437: Change to "...downstream of the sill."

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: Yes, a typo, which we will fix.

COMMENTS FROM REFEREE: Lines 477-478: This follows from the control limiting mainly the volume flux in the upper, active, layer.

AUTHORS' RESPONSE AND CHANGES IN MANUSCRIPT: Yes. We believe that we have explained this adequately and see no reason to modify the text.

# Internal hydraulic control in the Little Belt, Denmark. Observations of flow configurations and water mass formation.

Morten Holtegaard Nielsen<sup>1</sup>, Lars Chresten Lund-Hansen<sup>2</sup>, and Torben Vang<sup>2</sup>

<sup>1</sup>Marine Science & Consulting, Peder Lykkes Vej 8, 4. th., DK–2300 Copenhagen S, Denmark

5 <sup>2</sup>Marine Ecology, Biological Sciences, Aarhus University, Ole Worms Allé 1, Building 1135, DK–8000 Aarhus C, Denmark

*Correspondence to:* Morten Holtegaard Nielsen (mhn@msandc.dk)

**Abstract.** Internal hydraulic control, which occurs when stratified water masses are forced through an abrupt constriction, plays an enormous role in nature on both large and regional scale with respect to dynamics, circulation and water mass formation. Despite a growing literature on this subject surprisingly few direct observations have been made that conclusively show the existence of and the circumstances related to internal hydraulic control in nature. In this study we present observations from the Little Belt, Denmark, one of three narrow straits connecting the Baltic Sea and the North Sea. The observations, comprised of primarily along-strait, detailed transects of salinity and temperature, continuous observations of flow velocity, salinity and temperature on a permanent station and numerous vertical profiles of salinity, temperature, fluorescence and flow velocity in various locations, show that internal hydraulic control is a frequently occurring phenomenon in the Little Belt. The observations, which are limited to south-going flows of approximately two-layered water masses, show that internal hydraulic control may take either of two configurations, i.e. the lower or the upper layer being the active, accelerating one. This is connected to the depth of the pycnocline on the upstream side and the topography, which is both deepening and contracting toward the narrow part of the Little Belt. The existence of two possible flow configurations is known from theoretical and laboratory studies, but has never been observed in nature and reported before, we believe. The water masses formed by the intense mixing, which is tightly connected with the presence of control, may be found far downstream of the point of control. The observations show that these particular water masses are associated with chlorophyll concentrations that are considerably higher than in adjacent water masses, showing that control has a considerable influence on the primary production and hence the ecosystem in the area.

10  
15  
20

## 1 Introduction

25 In the recent years there has been a growing interest in the concept of internal hydraulic control, which occurs when a flowing, stratified fluid is subject to an abrupt constriction. In an idealized, layered flow this means that the Froude number, based on the speeds, depths and densities of the layers, exceeds unity at the constriction, implying that this acts as a control, determining the flow conditions in both upstream and downstream direction (e.g. Armi, 1986). This interest has often been stimulated by large scale phenomena such as the exchange of water masses between different ocean basins (e.g. Bryden et

30 al., 1994; Whitehead, 1998; Hansen et al., 2001; Girton and Sanford, 2003; Timmermans et al., 2005; Wilkenskjeld and  
Quadfasel, 2005), which is associated with transport of heat and salt and so may play an important role for the global  
thermohaline circulation (e.g. Clark et al., 2002; Vellinga and Wood, 2002). On a regional scale, internal hydraulic control  
has also attracted attention, for example in connection with estuarine circulation and bottom water renewal in semi-enclosed  
shelf seas and biological and environmental effects thereof (e.g. Nielsen, 2001; Roman et al., 2005; Hietala et al., 2007;  
35 Gregg and Pratt, 2010).

The purpose of this study is to present and discuss observations of internal hydraulic control (in the following and where  
ambiguities can be ruled out just referred to as hydraulic control) made in the Little Belt, Denmark (see Fig. 1). We find this  
pertinent for mainly two reasons. First, the conditions under which hydraulic control is occurring in nature are often very  
40 complex, including, primarily, irregular topography or stratification, large breadth-to-length and breadth-to-depth ratios,  
friction, mixing and the rotation of the earth (e.g. Farmer and Denton, 1985; Nielsen, 2001; Gregg and Pratt, 2010). In a  
great deal of existing theoretical or laboratory studies that have lead to much of the existing knowledge on this subject some  
or all of these conditions have often been avoided in order to make the problem tractable. Most of these existing studies have  
been reviewed by Garrett (2004), Ivey (2004), Lane-Serff (2004) and Pratt (2004), all in the same journal issue. Given the  
45 complex conditions in nature it is often hard to assess the applicability of various theories and approaches, including results  
of numerical models (e.g. Helfrich and Pratt, 2003). Attempts at validation of existing theories and models have included  
comparison of calculated total flow rates with those observed (Whitehead, 1998; Borenäs and Lundberg, 2004; Pratt, 2004).  
However, bulk numbers would seem inappropriate for such comparisons as they may easily conceal errors in the theory or  
model as well as inaccuracies in the observations (Garrett, 2004).

50  
Second, as detailed in the following, surprisingly few direct observations seem to have been made that conclusively show the  
existence and the circumstances related to the occurrence of hydraulic control in stratified water masses in nature. Of  
particular interest in this respect are the details at the point of control, which plays a crucial role by ultimately determining  
the conditions in either direction of the flow (e.g. Armi, 1986). Under conditions that are not fully ideal the location of the  
55 point of control has been shown to be influenced by, e.g., bottom friction and entrainment (Pratt, 1986; Gerdes et al., 2002;  
Nielsen et al., 2004). Existing observations of hydraulic control in a two-layered system (or with respect to the first internal  
mode) in a strait include Farmer and Smith (1980), Farmer and Denton (1985), Farmer and Armi (1999), Nielsen (2001),  
Inall et al. (2004), Klymak and Gregg (2004) and Staalstrøm et al. (2015). Other observations of control with respect to the  
first internal mode were made by Nash and Moum (2001), who studied the stratified flow over an isolated bank on the  
60 continental shelf. Direct observations of control of higher internal modes have also been made, although these are more  
subtle and play a lesser role with respect to dynamics, circulation and water mass formation (e.g. Farmer and Denton, 1985;  
Gregg and Klymak, 2014). On the other hand, a large number of observational studies have suggested the existence of  
hydraulic control in various locations without being able to observe the phenomenon directly (e.g. Farmer and Armi, 1988;

Armi and Farmer, 1988; Pratt et al., 2000; Fristedt et al., 2005; Girton et al., 2006; Gregg and Pratt, 2010). In some of these  
65 cases pronounced asymmetry and intense mixing around the assumed point of control are absent, suggesting that friction  
plays an important role in keeping the flow subcritical. In other cases the existence of a complex topography or an irregular  
stratification imply that the observations are very difficult to interpret from a hydraulic point of view.

Direct observations of hydraulic control under natural conditions may indeed hold the information that could be used to  
70 either validate or improve existing theories. Numerous investigations of hydraulic control reported in the literature have  
clearly shown a controversy regarding the principal question, namely how to determine criticality? Various approaches are  
based on energy considerations, calculation of the speed of long internal waves or the requirement that the flow remains  
well-behaved at the point of control (Farmer and Denton, 1985; Lawrence, 1993; Hogg et al., 2001). See Garrett (2004) and  
75 references therein for a discussion of some of these issues. It seems that the various theoretical approaches lead to the same  
result only when the most restrictive and simplifying assumptions are applied. It is beyond the scope of the present study to  
pursue this issue. However, we do hope that the observations that we present here may stimulate new thoughts that could  
lead toward a unifying solution.

An important aspect of the Little Belt is that the topography is subject to relatively smooth variations in terms of both  
80 breadth and depth, in contrast to most of the other situations in nature in which internal hydraulic control has been observed  
directly (Farmer and Smith, 1980; Farmer and Denton, 1985; Farmer and Armi, 1999; Nash and Moum, 2001; Inall et al.,  
2004; Klymak and Gregg, 2004; Staalstrøm et al., 2015). This implies that the Little Belt may be compared with and  
understood in terms of some simplified, existing theoretical and laboratory studies, such as Armi (1986) and Lawrence  
(1993). For the same reason we have limited ourselves to situations in which the stratification can be approximated by two  
85 layers and where the barotropic forcing is relatively strong, driving both layers in the same direction through the topographic  
constraint. This points to an important property of the role played by hydraulically controlled flows in nature, i.e. that the  
control is determining the ratio between the flows in the two layers (in a two-fluid system) whereas the total flow is  
determined by the barotropic forcing and, typically, the total friction in the system. An important finding by Armi (1986) is  
90 that in case of a two-layer flow through a contraction either of the two layers may become the active, accelerating one at the  
control. In Armi's laboratory experiment the solution was determined by selective withdrawal of the water masses on the  
downstream side of the contraction. In nature, however, the solution is much more likely to be determined by the total flow  
and the interface depth on the upstream side of the contraction. To our knowledge, observations in nature showing such two  
possible flow configurations in the same location have never been reported.

95 ~~Downstream of the point of control, where the flow is supercritical, intense mixing is taking place, resulting from growing  
instabilities at the pycnocline (e.g. Pawlak and Armi, 2000). This mixing, which has been found to be an order of magnitude  
more efficient than in the subcritical part of the flow (Prastowo et al., 2009, and references therein), is centered around the~~

100 pycnocline and tends to break down the stratification completely. As a result of the intense mixing associated with hydraulic control new water masses are formed that are distinctly different from those on the upstream side of the point of control. In supercritical flows the efficiency of mixing between the water masses is an order of magnitude higher than in subcritical flows (Prastowo et al., 2009, and references therein). This is due to growing instabilities at the pycnocline (Pawlak and Armi, 2000). This implies that downstream of the point of hydraulic control intense mixing is taking place, causing the formation of new water masses that are distinctly different from those on the upstream side.  conditions at the point of control have a strong bearing on the properties of these new water masses. For reasons of continuity, if the mixing in the supercritical part of the flow is strong enough to make the water masses completely homogeneous, the salinity on the downstream side is directly determined by the salinities of the water masses on the upstream side and the ratio of the flow rates at the control (e.g. Arneborg, 2016).  large scale it is well-known how through or overflows at prominent topographic constraints, e.g. the Denmark Strait or the Gibraltar, lead to the formation of water masses that may influence the circulation and can be traced far away from their locations of origin (Price et al., 1993; Dickson and Brown, 1994; Price and Baringer, 1994; Girton and Sanford, 2003). On a small or regional scale it is of interest to find out to what extent localized physical processes, especially energetic and efficient processes such as hydraulic control, are able to contribute to the formation of water masses and drive other processes (Gargett et al., 2003; Gregg and Pratt, 2010). Previously we have suggested that hydraulic control in the Little Belt could be accounting for enhanced primary production and oxygen depletion near the bottom, observed in wide areas in either direction away from the narrow part (Lund-Hansen et al., 2008). This question is explored further here.

115

In the next section a general description of the Little Belt is given. Then we explain the methods that were used to make the various observations, consisting of along-strait transects of temperature and salinity and vertical profiles and point measurements of temperature, salinity, fluorescence and/or current velocity from a number of locations. Then we present the observations and discuss their significance in relation to existing theoretical and laboratory studies. Then we provide a more general discussion of the findings, and finally conclusions are given.

120

## 2 Physical setting

The Little Belt is one of the three straits that form an important part of the transition area between the Baltic Sea and the North Sea (see Fig. 1). The Little Belt is comparatively long and narrow, its most prominent part being Snævringen, a 20 km long section in which the width decreases to about 1 km. At either ends of Snævringen, named Tragten and Bredningen, the changes of cross-sectional area are considerable, the widths reducing by 5 km or more over distances of about 2 km. The narrow part is relatively deep, the depth ranging from 30 to 50 m, whereas in the contracting parts to the north and to the south the depths are about 20 m or less. The water masses being forced through the narrow part of the Little Belt are thus experiencing both a reduced width and an increasing depth.

125

130 The currents through the Little Belt and the other Danish straits are much influenced by the meteorological conditions, which  
determine the water level variations in the North Sea and the Kattegat on the one side and in the Baltic Sea on the other  
(Jakobsen et al., 1997; Rohde, 1998). These result in flows into or out of the Baltic Sea of up to several days of duration,  
during which the water masses at the upstream end are advected through the straits. For instance, the water masses in the  
Kattegat are normally strongly stratified with salinities ranging from about 15 to 34 (Rasmussen, 1997; Nielsen, 2005), and  
135 in connection with inflow to the Baltic Sea these relatively quickly reach the narrow part of the Little Belt. In the inner  
Danish waters tidal waves are generally much smaller than the meteorologically induced water level variations, and in the  
Baltic Sea tides are practically absent (Jakobsen, 1980). However, in some places, one of which is the narrow part of the  
Little Belt, tidal flows of considerable magnitude can be found (Jakobsen and Ottavi, 1997). We expect this to be due to  
funneling and amplification of the tidal waves in the contracting area to the north of the narrow part of the Little Belt. Given  
140 typical flow speeds of  $0.5 - 1.0 \text{ m s}^{-1}$  the water masses are advected through the narrow part of the Little Belt in less than 12  
hours. Away from the narrow part the water masses are subject to much slower advection.

Because of the little significance for the exchange of the Baltic Sea, the Little Belt has received little attention in the  
scientific literature, see primarily Jakobsen and Ottavi (1997) and Lund-Hansen and Vang (2004). Other descriptions of the  
145 conditions in the areas between the Baltic Sea and the North Sea may be found in Stigebrandt (1983), Gustafsson (2000),  
Nielsen (2001) and references therein. Attention to the exchange through the three straits has often been motivated by  
inflows of dense water masses, which increase the salinity and relieve anoxic conditions in the deep parts of the Baltic Sea  
(e.g. Lass and Mohrholz, 2003). In this respect, the Little Belt normally plays a small role (Jakobsen, 1995). Other aspects  
related to the present study are described and discussed in Kepp et al. (2006) and Lund-Hansen et al. (2008).

### 150 **3 Methods**

This study is based on a part of a large set of observations made in and around the narrow part of the Little Belt (Fig. 1). The  
observations that we are using here were made on five separate occasions during April, May and June 2004. On all occasions  
south-going flow was taking place, forcing the strongly stratified water masses of the Kattegat through the Little Belt. The  
timings of the observations (except for the observations shown in Fig. 2) are indicated on the time axes in Fig. 3. The  
155 observations were made in the following four ways. 1) Transects of salinity, temperature and density along and across the  
strait were made using a ScanFish from EIVA, an undulating platform which was towed behind a ship. The ScanFish was  
equipped with an AROP 1000 CTD (Conductivity, Temperature, Depth) from Geological & Marine Instrumentation (GMI).  
Here and elsewhere we are using the Practical Salinity Scale, and density was calculated from salinity and temperature  
according to UNESCO (1981). The undulating speed of the ScanFish implied that a profile from the surface to the bottom  
160 was made in a horizontal distance of roughly 1 km. 2) At a permanent station located at the Old Bridge (Fig. 1) velocity was  
measured using a 300 kHz Workhorse Sentinel ADCP (Acoustic Doppler Current Profiler) from Teledyne RD Instruments

(RDI), which was placed on the seabed looking upward. The vertical velocity profile, recorded every 30 minutes, was determined using 150 pings and a bin depth of 1 m. Salinity and temperature, also recorded every 30 minutes, were measured at five, fixed depths using a Coastlog system from GMI. Here we are using data from the depths of 5 m (the uppermost sensor), 15 m (the middle sensor) and 22 m (the lowermost sensor). The total depth at the station was about 37 m. 3) At a few stations in the northern part of the Little Belt vertical profiles of salinity, temperature, density and velocity were observed from a ship. The profiles of salinity, temperature and density were observed using an AROP 2000 CTD from GMI. The velocity profiles were observed using a 600 kHz Workhorse Monitor ADCP from RDI, which was mounted over the side of the vessel looking downward, and calculated from about 100 pings and a bin depth of 0.5 m. 4) At a number of stations throughout the Little Belt vertical profiles of salinity, temperature, density and fluorescence were observed from a ship using an AROP 1000 CTD from GMI equipped with a standard fluorometer. The fluorescence data were converted to concentration of chlorophyll a based on a large number of filtered water samples in which the chlorophyll a content was determined by spectrophotometry. Other parts of the entire data set have been used and discussed by Kepp et al. (2006) and Lund-Hansen et al. (2008).

#### 4 Observations and discussion

Figure 2 shows a 50 km long transect of density along much of the Little Belt, covering the narrow part (roughly from 17 to 37 km on the length axis) as well as large parts of the contracting parts at either end. The observations were made using the ScanFish along the longer track as indicated in Fig. 1. The observations were carried out on 14 April 2004 between 09:10 and 15:07 (all times reported here are UTC), during a week-long period of weakly southward flow on the average, but with tidal flows of considerable magnitude, reaching an amplitude of about  $1 \text{ m s}^{-1}$  at the Old Bridge (data not shown). The transect shows a typical example of the conditions found in the Little Belt. At the northern end the water masses resemble those typically found in the Kattegat to the north, being strongly stratified and approximately two-layered, which may be attributed to a strong degree of wind mixing (Nielsen, 2005). In this case the water mass properties are ranging from a salinity and a temperature of about 18.5 and 6.7 °C, respectively, in the upper layer to about 26 and 4.6 °C in the lower layer, cf. the salinity-temperature plot shown in panel (b) in Fig. 2. At the southern end the water masses are only weakly stratified, and distinct layers are absent. In this case the water mass properties range from a salinity and a temperature of about 17 and 7.0 °C at the surface to 19 and 6.0 °C toward the bottom, cf. panel (d) in Fig. 2, i.e. of a different origin than the water masses found to the north. We attribute the weak and linear stratification in the southern part of the Little Belt to the depth being very variable, causing uneven mixing and transport due to wind and frequent restratification to take place. Typically, the water masses found in the narrow part of the Little Belt are well-mixed in the vertical, in this case showing little variation around a salinity and a temperature of 20.0 and 6.0 °C, cf. panel (c) in Fig. 2. In the present situation the well-mixed water masses found in the narrow part of the Little Belt are thus the result of mixing between the two water masses found to the north, the mixing ratio being roughly 80 % upper layer water and 20 % lower layer water.;

195 During the period in which the observations shown in Fig. 2 were made there was a weak inflow to the Baltic Sea, which we also refer to as southward flow, of the strongly layered and rather saline water masses of the Kattegat. At the same time, however, the tidal currents in the narrow part were strong enough to advect the water masses back and forth. From the transect in Fig. 2, which was carried out during a peak of southward flow (data not shown), it is apparent that the two-layered structure to the north is undergoing rapid adjustment as the water masses are forced toward the narrow part of the Little Belt. Furthermore, we note that the pycnocline between the water masses to the north is located deeply and that the lower layer seems to be active and accelerating as the water masses approach the narrow part. Under these circumstances the salinity in the well-mixed water masses on the downstream side indicates that the hydraulic control allows for an inflow of 80 % upper layer water and 20 % lower layer water from the upstream side. Given the deep location of the pycnocline, this would seem a probable result. We have previously argued that this rapid adjustment is connected with hydraulic control and that the intense mixing of the water masses is a result thereof (Lund-Hansen et al., 2008). The observations shown in Fig. 2 are not providing new information in this respect. However, the observations to be shown in the following will shed new light on hydraulic control in this area, including details of the adjustment and the effects on the formation of water masses. What Figs. 1 and 2 do show clearly though is that when well-mixed water masses are exported from the deep, narrow part of the Little Belt, covering a depth of about 50 m and a width of about 1 km, they are brought upward in the water column and are able to spread over a wide, shallow area, less than 20 m deep and several kilometres wide. Even though the mixed water masses could still be confined relatively deeply in the water column, this upwelling could well drive an increased growth of phytoplankton (Lyngsgaard et al., 2014).

Figure 3 shows time series of velocity, salinity and temperature obtained at the fixed station at the Old Bridge (see Fig. 1). The time series cover the two periods that include the observations shown in Figs. 4 through 8, the timings of these observations being indicated. The observations show the conditions that often prevail in the narrow part of the Little Belt. There is a strong tidal signal of diurnal frequency and amplitude ranging from the 0.5 to 1 m s<sup>-1</sup>. In addition, there is a flow component that varies on a longer time scale, of a few days or so, which is primarily caused by wind-generated water level variations in the Kattegat and in the western Baltic Sea (Jacobsen, 1980). This flow is of a considerable magnitude as well and may reach values of 1 m s<sup>-1</sup> or more. Figure 3 also shows that the water masses in the narrow part of the Little Belt are usually weakly stratified, despite the strong stratification often found at least in the Kattegat to the north (Nielsen, 2005). Given the magnitude of the barotropic flow we can quickly deduce that the flow in the narrow part of the Little Belt is often supercritical with respect to all internal modes. This circumstance was also pointed out by Jakobsen and Ottavi (1997). This is so even for the strongest stratification that one can imagine, which is a two-layer structure corresponding to the upper and lower layer water masses of the Kattegat of a density difference of about 14 kg m<sup>-3</sup> (Nielsen, 2005). In a 30 m water column of layers of equal thickness this corresponds to a long-wave speed of about 1 m s<sup>-1</sup> (e.g., Armi, 1986). As the cross-sectional area increases in either direction we can also deduce that situations of supercritical flow in the narrow part are associated

with subcritical flow on the upstream side and so that the flow must often be subject to hydraulic control, the control point to be found at the upstream contracting part of the Little Belt. We may therefore conclude that our assumptions so far about the observations of hydraulically controlled flows have been correct (Lund-Hansen et al., 2008). Further, we may point out that this finding is significantly above measurement errors. Thus, various effects that could dampen the flow below criticality, such as friction or mixing, are seen to be of less importance (Pratt, 1986; Nielsen et al., 2004). On the other hand, hydraulic control in the Little Belt is an intermittent phenomenon that is limited to parts of the tidal cycle if the barotropic flow is weak, which is the case in some of the periods shown in Fig. 3. The length scales of the contracting parts, about 2 km, are short compared to the distance travelled by a baroclinic wave during a tidal period (more than 10 km). This implies that the hydraulic adjustment is well toward the quasi-steady limit (Helfrich, 1995).

Another along-strait transect, obtained using the ScanFish as described above, is shown in Fig. 4. It covers the contracting part to the north and some of the narrow part, as shown by the line and circles in the detailed inset in Fig. 1. The transect was made on 16 June 2004 from 12:22 to 13:37 at the time of which the flow was southward through the Little Belt, the depth-averaged flow being well above  $1 \text{ m s}^{-1}$  for a period of about 24 hours, cf. Fig. 3. At the upstream end of the transect the water column is approximately two-layered, the salinity and the temperature being about 22 and  $13.5 \text{ }^\circ\text{C}$  in the upper layer and 29 and  $7.5 \text{ }^\circ\text{C}$  in the lower layer, cf. panel (b) in Fig. 4. Toward the narrow part of the Little Belt the water column is undergoing rapid adjustment, the pycnocline dropping toward the bottom from a depth of less than 10 m over a distance of a few kilometres, and the lower layer being subject to rapid dilution with upper layer water, cf. Figure 4, panel (c). In the narrow part of the strait the water masses are almost well-mixed, the mixing ratio being around 60 % upper layer water and 40 % lower layer water compared to the upstream side, cf. Figure 4, panel (d). From these circumstances we may safely conclude that the flow is subject to hydraulic control and that the lower layer is the active one. Furthermore, since the water column becomes fully mixed on the downstream side we may conclude that the hydraulic control allows a ratio of flow rates of the inflowing water masses equal to the mixing ratio. Due to shallower depth of the pycnocline in this situation compared to the situation shown in Fig. 2, a lower layer flow of 40 % of the total flow seems probable. Without accompanying observations of flow velocity we cannot say where the point of control is located though. Similar observations made some hours earlier on the same day along roughly the same track show the same picture and indicate the same location of the point of control (Lund-Hansen et al., 2008).

Just before the along-strait transect in Fig. 4 was made the two cross-strait transects of density shown in Fig. 5 were carried out. These cross-strait transects confirm that the water column is undergoing considerable adjustment as the narrow part of the Little Belt is approached, i.e. the interface dropping 5 – 10 m and mixing taking place. Both transects show some density variation across the strait indicating that geostrophy plays a role, which could make it difficult to interpret the density structure in the along-strait transect. However, since the Rossby radius of deformation is of the order of 10 km, rotational effects are relatively small in the part of the Little Belt concerned with hydraulic control. We note though that as the density

stratification is broken down in the supercritical part of the flow, rotational effects as well as curvature could become much more important. This is discussed more below. The two cross-strait transects along with the map in Fig. 1 show some important details of the topography. One is that following the deep, central part there is little change of width, whereas the bottom depth is increasing considerably. Another is that following the upper shallow part of the strait width is decreasing considerably as the narrow part of the Little Belt is approached. In a situation of two layers as considered here we may thus assume the following. In case of a relatively deep pycnocline, as is the case in Figs. 4 and 5, the lower layer will primarily experience a drop of the bottom, suggesting that the lower layer becomes the active one and the flow resembling that of 265 | spilling down a slope (e.g., Pawlak and Armi, 2000). In that case one would expect to find the point of control near the top of the slope. On the other hand, in case of a relatively shallow pycnocline the upper layer will be much influenced by the decreasing width. This suggests that the upper layer becomes the active one, that the flow resembles that through a contraction and that the point of control is to be found near the narrowest part of the contraction (e.g. Armi, 1986). In nature, hydraulic control are also influenced by other circumstances, e.g., friction or mixing, both of which tend to move the control point in the downstream direction (Pratt, 1986; Nielsen et al., 2004).

275

The vertical profiles of density and velocity shown in Fig. 6 were recorded in a situation much like the one in Figs. 4 and 5. The observations were obtained on 11 May around 19:00 during which the flow was southward of a speed of about 1 m s<sup>-1</sup> at the Old Bridge, cf. Fig. 3. These observations were made from a ship using a CTD that was profiled vertically through the water column and an ADCP that was mounted over the side looking downward. The profiles were recorded at two stations 280 | near what is assumed to be the location of the control point. The observations show a roughly two-layered water column with a density difference of about 8 kg m<sup>-3</sup>. At the upstream station the pycnocline was located at a depth of about 15 m, i.e. somewhat deeper than what was found in Figs. 4 and 5, and the velocity ranged from 0.4 to 0.6 m s<sup>-1</sup> with the upper layer moving slightly faster. At the other station, located about 2 km in the downstream direction, the pycnocline had dropped to a depth of about 30 m, the upper layer velocity was still around 0.6 m s<sup>-1</sup>, but the lower layer velocity had increased to about 285 | 1.2 m s<sup>-1</sup>. We note that there were significant secondary currents, which we attribute to the complexity of the topography as well as geostrophic adjustment.

The combination of observations of stratification and velocity in this situation allows us to calculate the composite Froude number. Since the relative density,  $\delta = (\rho_2 - \rho_1) / \rho_2$ , between the layers is small the composite Froude number can be 290 | approximated by

$$G^2 = F_1^2 + F_2^2,$$

where  $F_i^2 = U_i^2 / \delta g y_i$ ,  $y$  and  $U$  refer to the layer depth and velocity respectively, and indices 1 and 2 refer to the upper and the lower layer respectively (Armi, 1986). Taking layer depths of approximately 15 and 15 m at Station G53 and

30 and 15 m at Station G51, we find composite Froude numbers of  $G^2=0.31+0.14=0.45$  at G53 and  
295  $G^2=0.16+1.25=1.41$  at G51. As we have shown that the flow can be considered quasi-steady the increase of the  
composite Froude number to a value considerably above unity show that the flow is subject to hydraulic control and that in  
this situation, the lower layer being the active one and the flow spilling down the slope, the point of control is located  
somewhere between the two stations. This location is a bit downstream of the top of the slope, suggesting that friction and/or  
entrainment play a significant role in determining the control (Pratt, 1986; Nielsen et al., 2004).. This fact is not changed  
300 much if deviations from the two-layer assumption were to be taken into consideration, e.g. by calculating the effects of the  
density and the velocity profiles, cf. Garrett (2004). We note that the changes of layer depths and velocities are taking place  
rather abruptly, also indicative of the presence of control.

Figures 7 and 8 show a number of vertical profiles of density and chlorophyll a along with the corresponding distributions of  
305 salinity and temperature obtained on two separate occasions on stations covering most of the Little Belt, cf. Fig. 1. Both  
situations are concerned with inflow to the Baltic Sea, and in both cases the water masses on the upstream side are strongly  
stratified and approximately two-layered, as seen in the top rows of the two figures. An important difference between the two  
situations is related to the depth of the pycnocline on the upstream side. In the first situation the pycnocline is relatively  
deep, and in the other it is relatively shallow, which is seen to result in a difference in the hydraulic response between the two  
310 situations. The observations shown in Fig. 7 were made on 4 and 5 May 2004 during a weak southward flow on the average,  
but at the same time the tidal flows were considerable, the tidal amplitude being about  $0.5 \text{ m s}^{-1}$ , cf. Fig. 3. On the northern  
stations, shown in the upper panel in Fig. 7, the two water masses are characterized by a salinity and a temperature of about  
17 and  $10 \text{ }^\circ\text{C}$  in the upper layer and about 30 and  $5 \text{ }^\circ\text{C}$  in the lower layer, and the pycnocline is found at a depth of 12-15 m.  
In the narrow part of the Little Belt, seen in the middle row, the water masses are still mostly two-layered, and the upper  
315 layer is having roughly the same properties as found on the upstream side. However, in the lower layer a strong entrainment  
of upper layer water has taken place, resulting in a salinity and a temperature of about 24 and  $7 \text{ }^\circ\text{C}$  at Station 20. This  
corresponds to a mixing ratio of roughly 40 % upper layer water and 60 % lower layer water compared to the upstream side.  
At the same time the pycnocline has dropped to almost 20 m, strongly indicating that the flow is hydraulically controlled and  
that the lower layer is the active, accelerating one. At Station 21, south of the narrow part and seen in the lower row in Fig. 7,  
320 water masses of the same properties as produced by the strong entrainment in the narrow part are found, the mixing ratio  
having increased further to 80 % upper layer water and 20 % lower layer water compared to the upstream side. Much further  
to the south the density structure is almost the same, but the water masses are much different, showing that in this situation  
the mixed water masses resulting from hydraulic control in the narrow part are primarily found locally around the  
contraction and have not had time to be advected further southward.

325

The observations shown in Fig. 8 were made on 18 and 19 May 2004 during a period of moderate inflow to the Baltic Sea and tidal flows of an amplitude of about  $0.5 \text{ m s}^{-1}$ , which was not enough to reverse the overall flow (see Fig. 3). Notably, the pycnocline to the north is rather shallow, at a depth of 6-8 m, and the water masses are characterized by a salinity and a temperature of 18 and  $12 \text{ }^\circ\text{C}$  in the upper layer and about 30 and  $6 \text{ }^\circ\text{C}$  in the lower layer. In the narrow part of the Little Belt the lower layer is only modified slightly. But in the upper part of the water column a strong entrainment of lower layer water appear to have taken place., the mixed water masses consisting of up to about 80 % upper layer water and 20 % lower layer water compared to the upstream side. Furthermore, the density structure of the upper layer resembles that of a supercritical flow (e.g. Pawlak and Armi, 2000), all of which indicates that the flow is subject to hydraulic control and that the upper layer is the active, accelerating one in this situation. In fact, the observations made at the permanent station at the Old Bridge (Fig. 3), which is located about midway between Stations 51 and 20, show that the upper part of the water column is indeed moving considerably faster than both the average and the lower part. Thus, compared to the situation shown in Fig. 7 a different flow configuration is found. The cross-strait transects shown in Fig. 5 suggest that in case of a shallow pycnocline the upper layer is subject to an abrupt change of width when approaching the narrow part of the Little Belt, which is probably decisive for the flow configuration. Again, the water masses formed by hydraulic control are found locally downstream of the narrow part (at Station 21), whereas further to the south the water masses are much different. However, in this situation some of the water masses formed in the narrow part are found near the bottom at Station 27, some 30 km to the south of the narrow part, consisting of about 50 % upper layer water and 50 % lower layer water compared to the upstream side.

The CTD profiles shown in Figs. 7 and 8 were not accompanied by observations of flow velocity. Therefore we cannot provide direct evidence that these situations were subject to hydraulic control. However, with the south-going flow at the Old Bridge reaching peaks of almost  $1 \text{ m s}^{-1}$  and exceeding  $1.5 \text{ m s}^{-1}$  during the periods covered by Figs. 7 and 8, respectively, and the stratification being somewhat weaker than on the upstream side, we have strong reason to assume the flow in the narrow part of the Little Belt to be supercritical. This assumption is substantiated by the water column undergoing rapid adjustment and strong mixing taking place as the flow enters the narrow part of the Little Belt. Accordingly, the flow must be hydraulically controlled in these situations and the points of control be located in the contracting part to the north. Due to the lack of observations of flow velocity at the stations shown in Figs. 7 and 8 we are unable to narrow down where the points of control for the two flow configurations are located.

An interesting picture emerges when considering the distributions of chlorophyll a in relation to the different water masses found in the two situations shown in Figs. 7 and 8. First, it is noticed that away from the narrow part of the Little Belt the concentration of chlorophyll a is having a maximum at some depth that is coinciding with the density gradient, and above and below it is considerably lower. This is clearly seen in both the upper and the lower panels of Figs. 7 and 8, showing the conditions on the upstream side and on the downstream side of the narrow part of the Little Belt, respectively. Such

360 subsurface maxima are a prominent feature in areas that are strongly stratified and in which the growth of phytoplankton in the surface mixed layer is dependent on mixing across the pycnocline (e.g. Lyngsgaard et al., 2014). In contrast to this, at the stations where the water masses formed in connection with hydraulic control are found, both in the narrow part of the Little Belt (the middle panels) and on the downstream side (the lower panels), it is seen how these particular water masses are associated with markedly higher concentrations of chlorophyll a. The following two specific examples may help to clarify  
365 this mechanism. In Fig. 7 in the deep part of the water column at Station 21 mixed water masses containing up to roughly 20 % lower layer water and 80 % upper layer water from the upstream side are found, but the concentration of chlorophyll a is as high or even slightly higher than in the upper layer on the upstream side. In Fig. 8, also at Station 21 the mechanism is even clearer. First, the upper part of the water column consists of water masses containing roughly 60 % upper layer water from the upstream side whereas the chlorophyll concentration is well above that found in the upper layer on the upstream  
370 side. Second, in the lower part of the water column at Station 21, the content of upper layer from the upstream side is around 40 % whereas the chlorophyll a concentration is only slightly less than that found in the upper layer on the upstream side. Moreover, it is seen how these high concentrations are no longer confined to areas deep in the water column, but are extending toward the surface. Taking the increased chlorophyll a concentration as a proxy for primary production and increased phytoplankton concentration, it would seem that when the mixing associated with hydraulic control is bringing  
375 inorganic nutrients from the lower layer together with nutrient-depleted phytoplankton from the upper layer into the euphotic, upper part of the water column, the result is a quick response of the phytoplankton in terms of increased primary production and growth.

Previously, we have shown that the Little Belt is associated with locally high depth-integrated primary production and have  
380 suggested that this was closely related to mixing due to hydraulic control (Lund-Hansen et al., 2008). The observations in Figs. 7 and 8 clearly show that the water masses formed in connection with hydraulic control are indeed the ones associated with increased primary production and growth of phytoplankton. These water masses may be found locally around the narrow part of the Little Belt, but also some distance away. The latter is seen at Station 27 in Fig. 8, where the water masses formed in connection with hydraulic control are found near the bottom, due to a higher density than the water masses  
385 generally found in the southern part of the Little Belt. Despite their deep location where little light is available these water masses contain notably more chlorophyll a than at Station 29 further to the south. Therefore we may conclude that our suggestion was correct.

## 5 Discussion

In the present study we have limited our investigations in the Little Belt to a small number of situations of two-layered,  
390 southward flow. We have observed the existence of hydraulic control in these situations with certainty. In addition we have observed that the topography and the depth of the pycnocline play important roles in determining the flow configuration.

Despite the limited number of observations in this study, the magnitude of the barotropic flow along with the stratification found to the north or to the south show that hydraulic control in one form or another occurs on a regular basis. This should include control also of higher internal modes and, necessarily, so-called approach control (Farmer and Denton, 1985; Armi, 1986). Further, more detailed observations will have to shed light on these circumstances.

When contemplating the significance for a system consisting of two basins connected by a strait it is fundamental to realize the role that hydraulic control is playing. In oceanic or estuarine flows the total exchange of water masses through the strait is usually determined by the water level difference between the basins, which equal the head loss along the strait. In a two-layer flow the effect of hydraulic control is to determine at what proportion the total flow is divided between the flows of the different water masses. A good example is Øresund, the second largest of the three straits in the transition area between the Baltic Sea and the North Sea, in which the total flow is mainly determined by the frictional loss at the shallow and wide Drogden Sill (Jakobsen et al., 1997). In connection with southward flow hydraulic control in the contraction in the deep, northern part of Øresund is determining the proportion of upper and lower layer water of the Kattegat that can pass the Drogden Sill and flow into the Baltic Sea (Nielsen, 2001). Knowing the relation between the stratification and the flow at the point of control, i.e., a so-called weir formula (Pratt, 2004), one could ideally determine the variables of the flow as well as the dynamical properties of the system. In the Little Belt a similar functionality exists.

In the present study we have found that during periods of two-layered flow, which occurs often due to the stratification in the Kattegat (Nielsen, 2005), hydraulic control in the northern part of the Little Belt may be associated with two different flow configurations, i.e. the active, accelerating layer being either the upper or the lower one. Which of the two flow configurations actually occurs is a function of the depth of the pycnocline on the upstream side and probably also the total flow (Armi, 1986). This is so because the upper and the lower layer are primarily influenced by the reduction in width and the slope of the deep, central part, respectively. Different configurations of the flow through a contraction are known from the laboratory experiments by Armi (1986). In nature one should also expect that locations exist where different flow configurations may occur. However, we believe that this is the first time that this has been observed and reported in the scientific literature.

The observations have shown that the flow configuration seems to have a strong effect on the water masses that are being formed in connection with hydraulic control. When the lower layer is the active one, entrainment of upper layer water results in water masses of relatively low salinity. On the other hand, when the upper layer is the active one, entrainment of lower layer water leads to water masses that are relatively saline. In fact, the situation shown in Fig. 8 was associated with inflow to the Baltic Sea of the highest salinities in 2004 (data not shown). In addition, when considering the influence that hydraulic control might have on the conditions in the adjacent areas the amount of the formed water masses is important. This is a function of the ratio of flow in the two layers allowed by hydraulic control and the magnitude and the duration of the

barotropic flow. In the two situations shown in Figs. 7 and 8 the first was associated with a relatively weak barotropic flow and had had a short duration at the time of observation. In contrast, the second was both stronger and of longer duration and was associated with formation of water masses that were found far to the south of the narrow part of the Little Belt. Intermittency of the tidal flows makes the picture a little unclear, but, as mentioned previously, due to its relatively small  
430 volume the narrow part of the Little Belt is typically flushed in less than 12 hours and so is mostly reflecting the inflowing water masses at any given time.

Regardless of the flow configuration the observations reported here point to the fact that hydraulic control and the associated mixing in a strait is a mechanism that forms new, distinct water masses that may influence the circulation and other  
435 processes in the adjacent areas. In this case this is probably the reason why the Little Belt is generally associated with a higher primary production, higher concentrations of phytoplankton as well as recurrent oxygen depletion near the bottom (Jørgensen and Richardson, 1996; Lund-Hansen et al., 2008). Biological processes and the ecosystem dynamics as a whole are often driven by physical processes (Legendre and Demers, 1984). There may well be a great number of estuarine areas elsewhere in the world the functioning of which cannot be understood properly in the light of other prevailing physical  
440 processes, e.g. wind mixing, and where hydraulic control could play a decisive role (Roman et al., 2005). Again, knowing the relation between stratification and flow at the critical point, i.e. a so-called weir formula (Pratt, 2004), as well as the production of kinetic energy and a mixing efficiency (Prastowo et al., 2009), hydraulic control could potentially be used to quantify the transport of water masses and the mixing between them and to determine the influence on other processes. The controversy regarding how to determine criticality suggests that some comprehension and much work are needed before  
445 hydraulic control in nature can be utilized in this manner (Garrett, 2004). In fact, as pointed out by one anonymous referee, the mixing that we have observed in the Little Belt seems to be much more intense than in other observations of internal hydraulic control. This is possibly due to the relative long, narrow part of the Little Belt in which the flow may stay supercritical for a long distance. In comparison, most of the other examples of observations of hydraulic control known to us are concerned with a steep, relatively short sill with a dense, supercritical overflow spilling down the lee side of the sill  
450 (Farmer and Smith, 1980; Farmer and Denton, 1985; Farmer and Armi, 1999; Nash and Moum, 2001; Inall et al., 2004; Klymak and Gregg, 2004; Staalstrøm et al., 2015). In these examples mixing are confined to a relatively limited area downstream of the . The mechanisms of mixing under such circumstances are of considerable importance, but our observational material does not allow us to study these processes any deeper. Interested readers should consult, e.g., Klymak and Gregg (2004) and Staalstrøm et al. (2015).

455 As discussed above we expect the stratification in the southwestern Kattegat to have a strong influence on the effect that hydraulic control has on the inflow of water masses to the Baltic Sea. During the winter time the pycnocline is usually deeply located, due to strong wind mixing in the Kattegat (Rasmussen, 1997). In fact, the pycnocline may be found at such a depth that stratification is absent and that internal dynamics is not occurring in the Little Belt at all. During the summer, on

460 the other hand, the pycnocline may be relatively shallow, due to little wind mixing and heat input from the atmosphere. In addition, local winds may create up or downwelling. Over the short distance subject to internal hydraulic control and the hydraulic jump back to subcritical flow, winds only play a role in extreme situations. With respect to the observations that we have presented here the effect of wind mixing can be ruled out completely. In fact, only at Station 21 in Fig. 8, a bit downstream of the narrow part of the Little Belt, does it seem as could wind mixing have contributed to the mixing, seen as a  
465 homogeneous upper layer. The combination of baroclinic and barotropic forcing, which are varying somewhat independently, implies that hydraulic control in the Little Belt and its influence on the formation of water masses, inflow to the Baltic Sea, primary production and ecosystem dynamics is a complex and highly dynamic process. The full effect of this can probably only be examined and understood using an adequate model.

## 6 Conclusions

470 This observational study has shown that internal hydraulic control is a frequently occurring phenomenon in the Little Belt. The observations add to a surprisingly small number of studies that have conclusively shown the existence of control in stratified flows in nature. We have limited our observations to situations of south-going flow in which the water masses on the upstream side are approximately two-layered, facilitating a comparison with some existing laboratory and theoretical studies. As the layered water masses approach the narrow part of the Little Belt they are subject to both a deepening and a  
475 narrowing topography. The two layers undergo a rapid hydraulic adjustment with one of the layers accelerating strongly and intense mixing around the pycnocline taking place. In one situation vertical profiles of salinity, temperature and flow velocity show a considerable increase in the composite Froude number over a short distance of the flow. In this situation the lower layer is the active, accelerating one, the flow is spilling down the slope in the middle of the strait, and the point of control is found to lie a bit downstream from the top of the slope. In another situation in which the pycnocline on the  
480 upstream side is relatively shallow we observe the upper layer to be the active, accelerating one. In this situation the point of control is presumably located near the narrowest part of the upper, contracting part of the topography. Such two different flow configurations, which are known from theory and experiments in the laboratory, have never been observed in nature and reported before, we believe. Despite the limited number of observations the magnitude of the barotropic, partly tidally driven flows implies that internally critical flows are a recurring phenomenon under most circumstances in the Little Belt.  
485 Possibly this includes control of higher internal modes and approach-controlled flows.

The observations show that internal hydraulic control in the Little Belt is associated with intense mixing and formation of large amounts of water masses, which can be found far away on the downstream side of the control. These water masses are subject to much higher levels of chlorophyll *a* than the surrounding water masses, substantiating a previous study of ours that  
490 increased primary production in the adjacent areas is strongly influenced by control in the narrow part of the Little Belt. The two different flow configurations have a strong effect on the formation of water masses, the active, accelerating layer being

subject to intense entrainment and dilution. Thus, as the upper layer was observed to be the active one, water masses of the highest salinities in 2004 were produced and brought into the Baltic Sea.

### Author contributions

495 All authors took part in collecting the data. M. H. Nielsen carried out the data analysis and prepared the manuscript with contributions from the co-authors.

### Acknowledgements

500 The map in Fig. 1 is based on the Area Information System – Bathymetry for the Inner Danish Waters, copyright by the Danish Ministry of Environment. The observations shown in Figs. 2, 4 and 5 were made from R/V Tyra. The observations shown in Fig. 6 were made from R/V Genetica II. We are thankful to the crews of these vessels for their help and support. The data shown in Fig. 3 were obtained by the County of Vejle, and the data shown in Figs. 7 and 8 were obtained by the County of Funen. These data were made available through the Danish Natural Environment Portal. This study received some financial support from the Danish Natural Science Research Council (grant no. 4344-6757654). Finally, we would like to thank the three anonymous referees for their constructive criticism, with which we were able to improve the manuscript.

### 505 References

Armi, L.: The hydraulics of two flowing layers with different densities. *J. Fluid Mech.*, 163, 27–58, 1986.

Armi, L. and Farmer, D.: The flow of Mediterranean water through the Strait of Gibraltar. *Prog. Oceanogr.*, 21, 1–105, 1988.

[Arneborg, L.: Comment on “Influence of sea level rise on the dynamics of salt inflows in the Baltic Sea” by R. Hordoir, L.](#)

510 [Axell, U., Löptien, H., Dietze, and I. Kuznetsov. \*J. Geophys. Res. Oceans\*, 121, 2035–2040, doi:10.1002/2015JC011451, 2016.](#)

Borenäs, K. and Lundberg, P.: The Faroe-Bank Channel deep-water overflow. *Deep-Sea Res. II*, 51, 335–350, 2004.

Bryden, H. L., Candela, J. and Kinder, T. H.: Exchange through the Strait of Gibraltar. *Prog. Oceanogr.*, 33 (3), 201–248, 1994.

515 Clark, P. U., Pisias, N. G., Stocker, T. F. and Weaver, A. J.: The role of the thermohaline circulation in abrupt climate change. *Nature*, 415, 863–869, doi:10.1038/415863a, 2002.

Dickson, R. R. and Brown, J.: The production of North Atlantic Deep Water: Sources, rates and pathways. *J. Geophys. Res.*, 99 (C6), 12,319–12,341, 1994.

Farmer, D. and Armi, L.: The flow of Atlantic water through the Strait of Gibraltar. *Prog. Oceanogr.*, 21, 1–105, 1988.

- Farmer, D. and Armi, L.: Stratified flow over topography: the role of small-scale entrainment and mixing in flow establishment. *Proc. R. Soc. Lond. A*, 455, 3221–3258, 1999.
- Farmer, D. M. and Denton, R. A.: Hydraulic control of flow over the sill in Observatory Inlet. *J. Phys. Oceanogr.*, 90 (C5), 9051–9068, 1985.
- Farmer, D. M. and Smith, J. D.: Tidal interaction of stratified flow with a sill in Knight Inlet. *Deep-Sea Res.*, 27A, 239–254, 1980.
- 525 Fristedt, T., Lundberg, P. and Sigraý, P.: Technical note on possible hydraulic control of the exchange through the Oxdjupet passage in the Baltic Sea. *Estuar. Coast. Shelf Sci.*, 65, 526–534, 2005.
- Garrett, C.: Frictional processes in straits. *Deep-Sea Res. II*, 51, 393–410, 2004.
- Gargett, A. E., Stucchi, D. and Whitney, F.: Physical processes associated with high primary production in Saanich Inlet, British Columbia. *Estuar. Coast. Shelf Sci.*, 56, 1141–1156, 2003.
- 530 Gerdes, F., Garrett, C. and Farmer, D.: On internal hydraulics with entrainment. *J. Phys. Oceanogr.*, 32, 1106–1111, 2002.
- Girton, J. and Sanford, T.: Descent and modification of the overflow plume in the Denmark Strait. *J. Phys. Oceanogr.*, 33, 1351–1364, 2003.
- Girton, J. B., Pratt, L. J., Sutherland, D. A. and Price, J. F.: Is the Faroe Bank Channel overflow hydraulically controlled? *J. Phys. Oceanogr.*, 36, 2340–2349, 2006.
- 535 Gregg, M. C. and Klymak, J. M.: Mode-2 hydraulic control of flow over a small ridge on a continental shelf, *J. Geophys. Res. Oceans*, 119, 8093–8108, doi:10.1002/2014JC010043, 2014.
- Gregg, M. C. and Pratt, L. J.: Flow and hydraulics near the sill of Hood Canal, a strongly sheared, continuously stratified fjord. *J. Phys. Oceanogr.*, 40, 1087–1105. doi: 10.1175/2010JPO4312.1, 2010.
- Gustafsson, B.: Time-dependent modeling of the Baltic entrance area: 1. Quantification of circulation and residence times in the Kattegat and the straits of the Baltic Sill. *Estuaries*, 23, 231–252, 2000.
- 540 Hansen, B., Turrell, W. R. and Østerhus, S.: Decreasing overflow from the Nordic seas into the Atlantic Ocean through the Faroe Bank channel since 1950, *Nature*, 111, 927–929, 2001.
- Helfrich, K. R.: Time-dependent two-layer hydraulic exchange flows. *J. Phys. Oceanogr.*, 25, 359–373, 1995.
- Helfrich, K. R. and Pratt, L. J.: Rotating hydraulics and upstream basin circulation. *J. Phys. Oceanogr.*, 1651–1663, 2003.
- 545 Hietala, R., Lundberg, P. and Nilsson, J. A. U.: A note on the deep-water inflow to the Bothnian Sea. *J. Mar. Syst.*, 68, 255–264, 2007.
- Hogg, A. M., Winters, K. B. and Ivey, G. N.: Linear internal waves and the control of stratified exchange flows. *J. Fluid Mech.*, 447, 357–375, 2001.
- Inall, M., Cottier, F., Griffiths, C. and Rippeth, T.: Sill dynamics and energy transformation in a jet fjord. *Ocean Dynam.*, 54, 550 307–314, 2004.
- Ivey, G. N.: Stratification and mixing in sea straits. *Deep-Sea Res. II*, 441–453, 2004.

- Jacobsen, T. S.: Seawater exchange of the Baltic: measurements and methods. PhD dissertation. National Agency of Environmental Protection, Copenhagen, 1980.
- Jakobsen, F.: The major inflow to the Baltic Sea during January 1993. *J. Mar. Syst.*, 6, 227–240, 1995.
- 555 Jakobsen, F. and Ottavi, J.: Transport through the contraction area in the Little Belt. *Estuar., Coast. Shelf Sci.*, 45, 759–767, 1997.
- Jakobsen, F., Lintrup, M. J. and Møller, J. S.: Observations of the specific resistance in the Øresund. *Nord. Hydrol.*, 28, 217–232, 1997.
- Jørgensen, B.B. and Richardson, K. (Eds.): Eutrophication in coastal marine ecosystems. American Geophysical Union, 560 Washington, 281 pp., 1996.
- Kepp, R. R., Struve, A., Christiansen, C., Lund-Hansen, L. C., Nielsen, M. H. and Vang, T.: Transport and hydraulically-induced recycling of phosphorus in the North Sea-Baltic Sea transition zone. *Oceanologia*, 48 (2), 175–191, 2006.
- Klymak, J. M. and Gregg, M. C.: Tidally generated turbulence over the Knight Inlet sill. *J. Phys. Oceanogr.*, 34, 1135–1151, 2004.
- 565 Lane-Serff, G. F.: Topographic and boundary effects on steady and unsteady flow through straits. *Deep-Sea Res. II*, 51, 321–334, 2004.
- Lass, U. and Mohrholz, V.: On dynamics and mixing of inflowing saltwater in the Arkona Sea. *J. Phys. Oceanogr.*, 108 (C2), 3042, doi:10.1029/2002JC001465, 2003.
- Lawrence, G. A.: The hydraulics of steady two-layer flow over a fixed obstacle. *J. Fluid Mech.*, 254, 605–633, 1993.
- 570 Legendre, L. and Demers, S.: Towards dynamic biological oceanography and limnology. *Can. J. Fish. Aquat. Sci.*, 41, 2–19, 1984.
- Lund-Hansen, L. C. and Vang, T.: An inflow and intrusion event in the Little Belt at the North Sea–Baltic Sea transition and a related sub-surface bloom of *Pseudo-nitzschia pseudodelicatissima*. *Estuar. Coast. Shelf Sci.*, 59, 265–276, 2004.
- Lund-Hansen, L. C., Nielsen, M. H., Bruhn, A., Christiansen, C., Vang, T., Casado-Amezua, P., Richardson, K. and 575 Santaloria, L.: A consistent high primary production and chlorophyll-a maximum in a narrow strait – Effects of hydraulic control. *J. Mar. Syst.*, 74, 395–405, 2008.
- Lyngsgaard, M. M., Richardson, K., Markager, S., Nielsen, M. H., Olesen, M. and Christensen, J. P. A.: Deep primary production in coastal pelagic systems: importance for ecosystem functioning, *Mar. Ecol. Prog. Ser.*, 517, 15–33. doi:10.3354/meps11015, 2014.
- 580 Nash, J. D. and Moum, J. N.: Internal hydraulic flows on the continental shelf: high drag states over a small bank. *J. Geophys. Res.*, 106 (C3), 4593–4612, 2001.
- Nielsen, M. H.: Evidence for internal hydraulic control in the northern Øresund. *J. Geophys. Res.*, 106 (C7), 14,055–14,068, 2001.
- Nielsen, M. H.: The baroclinic surface currents in the Kattegat. *J. Mar. Syst.*, 55 (3–4), 97–121, 2005.

- 585 Nielsen, M. H., Pratt, L. J. and Helfrich, K. R.: Mixing and entrainment in hydraulically driven stratified sill flows. *J. Fluid Mech.*, 515, 415–443. doi:10.1017/S0022112004000576, 2004.
- Pawlak, G., and Armi, L.: Mixing and entrainment in developing stratified currents. *J. Fluid Mech.*, 424, 45–73, 2000.
- Prastowo, T., Griffiths, R. W., Hughes, G. O. and Hogg, A. M.: Effects of topography on the cumulative mixing efficiency in exchange flows, *J. Geophys. Res.*, 114, C08008, doi:10.1029/2008JC005152, 2009.
- 590 Pratt, L.: Hydraulic control of sill flow with bottom friction. *J. Phys. Oceanogr.*, 16, 1970–1980, 1986.
- Pratt, L. J.: Recent progress on understanding the effects of rotation in models of sea straits. *Deep-Sea Res. II*, 51, 351–369, 2004.
- Pratt, L. J., Deese, H. E., Murray, S. P. and Johns, W.: Continuous dynamical modes in straits having arbitrary cross sections, with applications to the Bab al Mandab. *J. Phys. Oceanogr.*, 2515–2534, 2000.
- 595 Price, J. F. and Baringer, M. O.: Outflows and deep water production by marginal seas. *Prog. Oceanogr.*, 33, 161–200, 1994.
- Price, J. F., Baringer, M. O., Lueck, R. G., Johnson, G. C., Ambar, I., Parrilla, G., Cantos, A., Kennelly, M. A. and Sanford, T. B.: Mediterranean outflow mixing and dynamics. *Science*, 259, 1277–1282, 1993.
- Rasmussen, B.: The near-surface horizontal buoyancy flux in a highly stratified region, Kattegat. *Estuar. Coast. Shelf Sci.*, 600 45, 405–414, 1997.
- Rohde, J.: The Baltic and North Seas: a process-oriented review of the physical oceanography. In: Robinson, A. R. and Brink, K. H. (Eds): *The Sea*, 11, 699–732, Wiley, London, 1998.
- Roman, M., Zhang, X., McGilliard, C. and Boicourt, W.: Seasonal and annual variability in the spatial patterns of plankton biomass in Chesapeake Bay. *Limnol. Oceanogr.*, 50 (2), 480–492, 2005.
- 605 Staalstrøm, A., Arneborg, L., Liljebldh, B. and Broström, G.: Observations of turbulence caused by a combination of tides and mean baroclinic flow over a fjord sill. *J. Phys. Oceanogr.*, 45, 355–368.
- Stigebrandt, A.: A model for the exchange of water and salt between the Baltic and the Skagerrak. *J. Phys. Oceanogr.*, 13, 411–427, 1983.
- UNESCO: International oceanographic tables, Vol. 3. UNESCO Technical Papers in Marine Science, 39. UNESCO, Paris, 610 1981.
- Whitehead, J. A.: Topographic control of oceanic flows in deep passages and straits. *Rev. Geophys.*, 36 (3), 423–440, 1998.
- Wilkenskjeld, S. and Quadfasel, D.: Response of the Greenland-Scotland overflow to changing deep water supply from the Arctic Mediterranean. *Geophys. Res. Lett.*, 32, L21607, doi:10.1029/2005GL024140, 2005.
- Timmermans, M.-L., Winsor, P. and Whitehead, J. A.: Deep-water flow over the Lomonosov Ridge in the Arctic Ocean. *J. Phys. Oceanogr.*, 35, 1489–1493, 2005.
- 615 Vellinga, M. and Wood, R. A.: Global climatic impacts of a collapse of the Atlantic thermohaline circulation. *Climatic Change*, 54, 251–267. doi:10.1023/A:1016168827653, 2002.

## Figure captions

620 Figure 1. Map of the Little Belt with insets showing the three straits connecting the Baltic Sea and the North Sea (upper left)  
and a detailed view of the narrow part and the contracting part to the north (right). The depth contours indicate 10 m (solid),  
20 m (dashed) and 30 m (dotted). The original place names that are briefly mentioned in the text are shown. The locations of  
the stations from which data have been used are shown (solid squares), OB implying the Old Bridge. The two along-strait  
survey lines, along which every 2 and 5 km are marked (circles), are where the density transects in Figs. 2 and 4 were made.  
The two cross-strait survey lines just north of the narrow part are where the density transects in Fig. 5 were made.

625

Figure 2. Along-strait transect of observations of density (minus  $1000 \text{ kg m}^{-3}$ ) given by contour lines (upper panel) and the  
distributions of salinity (in Practical Salinity Units) and temperature of the entire transect (panel a), at 5 – 10 km (panel b), at  
25 – 30 km (panel c) and at 45 – 50 km (panel d). In the salinity-temperature plots the lines are contours of constant density  
(minus  $1000 \text{ kg m}^{-3}$ ). The observations were made using a ScanFish (see text), the entire transect corresponding to more than  
630 40 undulations from surface to bottom. The transect was made on 14 April 2004 between 09:10 and 15:07 along the long  
survey line shown Fig. 1, the length of the transect being measured from the northern end.

Figure 3. Time series of observations from the station at the Old Bridge (see Figure 1) for 2 through 19 May and 14 through  
17 June 2004. The upper panel shows velocity measurements projected onto the main flow direction, positive values for  
635 northward flow through the Little Belt. The curves are the depth-averaged value (solid, thick), the bin box at a depth of about  
5 m (solid, thin), and the bin box at 3 m above the bottom (dashed, thin). The total depth at the station is about 37 m. The  
middle and the lower panel show the salinity (in Practical Salinity Units) and the temperature, respectively, measured at a  
depth of 5 m (solid, thin curve), 15 m (solid, thick curve) and 22 m (dashed, thin curve). The timings of the observations  
shown in Figs. 4 through 8 are indicated by the short, thick lines on the x axes, specified in the upper panel. (Some of the  
640 curves may be difficult to distinguish. Note though that the less saline and warmer water is found near the surface.)

Figure 4. Along-strait transect of observations of density (minus  $1000 \text{ kg m}^{-3}$ ) given by contour lines (upper panel) and the  
distributions of salinity (in Practical Salinity Units) and temperature of the entire transect (panel a), at 0 – 2 km (panel b), at  
7 – 9 km (panel c) and at 16 – 18 km (panel d). In the salinity-temperature plots the lines are contours of constant density  
645 (minus  $1000 \text{ kg m}^{-3}$ ). The observations were made using a ScanFish (see text), the entire transect corresponding to 15  
undulations. The transect was made on 16 June 2004 from 12:22 to 13:37 along the short survey line as shown Fig. 1.

Figure 5. Cross-strait transects of observations of density (minus  $1000 \text{ kg m}^{-3}$ ) given by contour lines. The transects were  
made using a ScanFish (see text) along the two cross-strait survey lines just north of the narrow part of the Little Belt as

650 shown in Fig. 1. The transects were made on 16 June 2004 from 09:45 to 10:56. The northern transect is shown in the left panel, the southern in the right, and in both cases the length axes are pointing from north-west to south-east.

Figure 6. Vertical profiles of density (thick lines) and velocity (thin lines) measured at the two stations G53 (left panel) and G51 (right panel), shown in Fig. 1. The observations were made on 11 May 2004 at 19:10 and 18:20 respectively. The  
655 velocity is projected onto the main flow direction (solid line; positive southward through the Little Belt) and perpendicular to this (dashed line; positive to the right of the main flow direction).

Figure 7. Vertical profiles of density (panels to the left) and chlorophyll a (panels in the middle), and the corresponding distributions of temperature and salinity (in Practical Salinity Units) (panels to the right). In the salinity-temperature plot the  
660 lines are contours of constant density (minus  $1000 \text{ kg m}^{-3}$ ). The observations were made on 4 May 2004 between 07:35 and 08:30 at stations 14 (circles), 16 (squares) and 18 (crosses), all shown in the upper row, on 4 May at 08:51 at station 51 (circles) and on 5 May at 05:17 at station 20 (squares), shown in the middle row, and on 5 May between 05:45 and 08:49 at stations 21 (circles), 27 (squares) and 29 (crosses), shown in the lower row.

665 Figure 8. Vertical profiles of density (panels to the left) and chlorophyll a (panels in the middle), and the corresponding distributions of temperature and salinity (in Practical Salinity Units) (panels to the right). In the salinity-temperature plot the lines are contours of constant density (minus  $1000 \text{ kg m}^{-3}$ ). The observations were made on 18 May 2004 between 08:19 and 09:16 at stations 14 (circles), 16 (squares) and 18 (crosses), all shown in the upper panel, on 18 May 09:34 at station 51 (circles) and on 19 May 05:26 at station 20 (squares), shown in the middle panel, and on 19 May between 05:52 and 08:41 at  
670 stations 21 (circles), 27 (squares) and 29 (crosses), shown in the lower row.