



Properties and dynamics of mesoscale-eddies in the Fram Strait from a comparison between two high-resolution ocean-sea ice models

Claudia Wekerle¹, Tore Hattermann^{2,3}, Qiang Wang¹, Laura Crews⁴, Wilken-Jon von Appen¹, and Sergey Danilov¹

¹Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany

²Norwegian Polar Institute, Tromsø, Norway

³Energy and Climate Group, Department of Physics and Technology, The Arctic University of Tromsø, Norway

⁴School of Oceanography and Applied Physics Laboratory, University of Washington, Seattle, USA

Correspondence: Claudia Wekerle (Claudia.Wekerle@awi.de)

Abstract. The Fram Strait, the deepest gateway to the Arctic Ocean, is strongly influenced by eddy dynamics. Here we analyse the output from two eddy-resolving models (ROMS and FESOM) with around 1 km mesh resolution in the Fram Strait, with focus on their representation of eddy properties and dynamics. A comparison with mooring observations shows that both models reasonably simulate hydrography and eddy kinetic energy. Despite differences in model formulation, they show relatively similar eddy properties. The eddies have a mean radius of 4.9 km and 5.6 km in ROMS and FESOM, respectively, with slightly more cyclones than anticyclones (ROMS: 54%, FESOM: 55%). The lifetime of detected eddies is relatively short in both simulations (ROMS: 10 days, FESOM: 11 days), and the mean travel distance is 35 km in both models. More anticyclones are trapped in deep depressions or move toward deep locations. The two models show comparable patterns of baroclinic and barotropic instability. However, ROMS has relatively stronger eddy intensity and baroclinic instability, possibly due to its smaller grid size and higher effective resolution. Overall, the relatively good agreement between the two models strengthens our confidence in their ability to realistically represent the Fram Strait ocean dynamics, and also highlights the need for very high mesh resolution.

1 Introduction

The Fram Strait, located between Svalbard and Greenland (Figure 1), is the deepest gateway that connects the Arctic Ocean and the North Atlantic via the Nordic Seas. Many important processes of climate relevance take place in this region. On the one hand, Atlantic Water (AW) carried northward by the West Spitsbergen Current (WSC, e.g. von Appen et al., 2016) enters the Arctic Ocean as its largest oceanic heat source. In the last decades, an increase in AW temperature has been observed in the Fram Strait, with implications for the Arctic Ocean's sea ice decline (Beszczynska-Möller et al., 2012; Polyakov et al., 2012). On the other hand, a part of the AW recirculates in the Fram Strait and continues southward in the East Greenland Current (EGC, e.g. de Steur et al., 2009). This water mass, which was densified on its way north to Fram Strait, contributes to the Denmark Strait overflow, which forms the dense part of the North Atlantic Deep Water, a key component of the Atlantic



meridional overturning circulation (Eldevik et al., 2009). Furthermore, cold and fresh Polar Water (PW) carried southward by the EGC is injected into the cyclonic Greenland Sea Gyre, impacting convection there (Rudels, 1995), and thus also the overflow across the Greenland Scotland Ridge.

25 The oceanic conditions in the Fram Strait are strongly turbulent. Already in the 1980's it was revealed by measurement campaigns such as the Marginal Ice Zone Experiments that eddies are abundant there (Johannessen et al., 1987b; Smith et al., 1984). They play an important role in shaping the ocean circulation and hydrography, sea ice and ecosystem:

(1) Some eddies are shed from the WSC and travel westward, driving the recirculation of warm and salty AW. This was shown by mooring measurements (Schauer et al., 2004; von Appen et al., 2016) and model simulations (Hattermann et al.,
30 2016; Wekerle et al., 2017), which revealed high levels of eddy kinetic energy (EKE) in the WSC and along the recirculation pathway.

(2) As AW recirculates, it subducts underneath cold and fresh Polar Water (PW) carried by the East Greenland Current (EGC). As shown by Hattermann et al. (2016), baroclinic instability leads to vertical eddy fluxes which may subduct AW.

(3) Once the Return Atlantic Water (RAW) crosses (likely eddy mediated) the Northeast Greenland continental shelf break,
35 part of it travels through a trough system towards the Northeast Greenland glaciers (Schaffer et al., 2017). An increase in its temperature might lead to the glaciers' destabilisation (Wilson et al., 2017), and it has been shown that eddy overturning is important for lifting AW onto the continental shelf in the Fram Strait (Tverberg and Nøst, 2009; Cherian and Brink, 2018).

(4) The marginal ice zone is shaped by eddies (Johannessen et al., 1987b) and submesoscale features (von Appen et al.,
40 2018). By means of idealised model experiments, Manucharyan and Thompson (2017) showed that cyclonic eddies can trap sea ice and carry it to warm waters, leading to enhanced melting rates.

(5) Eddy and filamentary structures are important features for the marine ecosystem. Among other effects, they play an important role in transporting nutrients into the euphotic zone for phytoplankton production, and can cause stratification within days, thereby increasing light exposure for phytoplankton trapped close to the surface (Mahadevan, 2016).

Eddies can be generated through both baroclinic and barotropic instabilities. In the presence of horizontal density gradients
45 and baroclinic instability, mesoscale eddies develop through the conversion of the available potential energy (APE) to EKE. Barotropic instability in contrast is associated with horizontal shear in jet-like currents, and eddies can be formed by receiving kinetic energy from the mean flow (Teigen et al., 2011). Eddies can also be steered or trapped by topography (Johannessen et al., 1987a). This steering modulates the conversion between eddy and mean kinetic energy, which can be directed in both ways. Fram Strait, featured with its complex topography, strong lateral gradients in temperature and salinity (warm and saline AW
50 in the eastern part, cold and fresh PW in the western part) and thus steep isopycnal slopes across the strait, strong convective events in the winter months and strong boundary currents (WSC and EGC), is thus a highly active and interesting region for studying eddy dynamics.

The Rossby radius of deformation, which characterises the spatial scale of eddies, is small in the Fram Strait, around 4–6 km in summer and 3–4 km in winter (von Appen et al., 2016). This poses problems for ocean models which typically operate
55 on coarser grids. Recently, high resolution ocean models focused on the Fram Strait region have emerged, which perform well in reproducing the observed eddy activity (Kawasaki and Hasumi, 2016; Hattermann et al., 2016; Wekerle et al., 2017).



Given the possible sensitivity of simulations to model numerics, to the complex bottom topography and ocean currents in the Fram Strait, it is not known whether the above cited models have a broad agreement on the representation of eddy dynamics in terms of eddy generation and propagation. Answering this question will not only add credence to our understanding of eddy dynamics, but also create a reference for developing parameterisations required by coarse resolution ocean models. The aim of this study is two-fold. First, we compare the output of two high-resolution, eddy-resolving ocean-sea ice models to answer the above question. We will show that there is good agreement in energy conversion that maintains eddy dynamics and in simulated eddy statistics as well, despite the fact that these models, namely ROMS (Hattermann et al., 2016) and FESOM (Wekerle et al., 2017), differ in many aspects such as numerical discretisation, horizontal and vertical mesh resolution, parameterisations, global vs. regional configurations. Second, we explore and describe the properties of eddies in the Fram Strait. We use an eddy-following approach to generate regional statistics focusing on the following questions: How are eddies spatially distributed? Are anticyclones or cyclones dominating? What is their typical size, lifetime and what are their main travel pathways?

2 Methods

2.1 Model description FESOM

Model output from the Finite-Element Sea-ice Ocean Model (FESOM) version 1.4 (Wang et al., 2014; Danilov et al., 2015) is used for eddy detection and tracking in this study. FESOM is an ocean-sea ice model which solves the hydrostatic primitive equations in the Boussinesq approximation and is discretised with the finite element method (Wang et al., 2008). In the vertical, z-levels are used. We use a global FESOM configuration that was optimised for the Fram Strait with regional resolution (grid size) refined to 1 km in this area, and a coarser resolution elsewhere (1° resolution throughout most of the world's oceans, 24 km resolution north of 40°N and 4.5 km resolution in the Nordic Seas and Arctic Ocean; Wekerle et al. (2017)). By comparing with the local Rossby radius of deformation (around 3–6 km in the Fram Strait, see above), this configuration can be considered as "eddy-resolving". It is forced with atmospheric reanalysis data from COREv.2 (Large and Yeager, 2008), and river runoff is taken from the interannual monthly data set provided by Dai et al. (2009). Tides are not taken into account in the FESOM configuration used here. The simulation covers the time period 2000–2009, and has daily output. In this study, we analyse model output for the years 2006–2009.

2.2 Model description ROMS

The second high-resolution model simulation used in this study is based on the Regional Ocean Modeling System (ROMS) (Budgell, 2005; Haidvogel et al., 2008; Shchepetkin and McWilliams, 2005, 2009) with a configuration optimised for Fram Strait and the waters around Svalbard (called S800). With 800 m x 800 m horizontal resolution, S800 is eddy resolving in Fram Strait. S800 was initialised with and forced at the ocean boundaries with daily ocean and sea ice data from a 4 km resolution pan-Arctic model called A4, together with tidal elevations from global TPXO tidal model (Egbert and Erofeeva, 2002). A4's initial state and boundary conditions were taken from monthly-averaged global reanalyses (Storkey et al., 2010).



Atmospheric forcing in A4 and S800 used 6-hourly ERA-Interim reanalysis (Dee et al., 2011). A4 was initialised in 1993, and following A4 spin-up S800 was initialised in January 2005. Analyses in this paper are done for the period of 2006–2009. Model characteristics of ROMS, and also of FESOM, are summarised in Table 1. Additional information about S800, including discussions of its ability to reproduce boundary current observations in Fram Strait and along the continental slope north of Svalbard, is given in Hattermann et al. (2016), Sundfjord et al. (2017), Crews et al. (2018) and Crews et al. (2019).

2.3 Eddy detection and tracking

Eddy detection and tracking algorithms are important tools to understand eddy properties such as their size, strength, lifetime and travel pathways. For datasets as large as the output of ocean models, automated methods need to be used. Eddy detection methods can be assigned to two categories, based either on (1) geometrical or on (2) physical characteristics of the flow field, or on a combination of both. In this study, we apply a method developed by Nencioli et al. (2010) to detect and track eddies simulated with ROMS and FESOM, which is based on the geometry of velocity vectors and thus belongs to the first category of methods. The eddy detection is based on four constraints derived from the general characteristics of velocity fields in the presence of eddies, e.g. the reversal of the flow field across the eddy centre. Two parameters, a and b , which determine the minimum size of detectable vortices, have to be set in the algorithm. After some sensitivity tests, we set $a = 4$ and $b = 3$, which equals the values used in the test case of Nencioli et al. (2010). Note that our mesh resolutions (800 m and 1 km in ROMS and FESOM, respectively) are similar to theirs (1 km). Eddy boundaries around each detected centre are determined by the outermost closed contour of the stream function field.

To cross-validate our results, we also the Okubo-Weiss criterion, which belongs to the second category of methods (Okubo, 1970; Weiss, 1991). Eddies are identified as areas where vorticity dominates over strain. More precisely, the area where the Okubo-Weiss parameter

$$OW = \underbrace{(\partial_x u - \partial_y v)^2 + (\partial_x v + \partial_y u)^2}_{\text{normal and shear component of strain}} - \underbrace{(\partial_x v - \partial_y u)^2}_{\text{relative vorticity}} \quad (1)$$

is below a threshold of $OW_0 = -0.2\sigma_{OW}$ with same sign of vorticity, where σ_{OW} is the spatial standard deviation of OW, is considered as an eddy (Isern-Fontanet et al., 2006). Here (u, v) is the horizontal velocity field, and f is the Coriolis parameter.

After eddies are detected, eddy tracks are computed by comparing eddy centres in successive time steps. More precisely, if two eddies at successive time steps lie within a search radius and have the same sense of rotation, they form a track. The eddy tracking scheme is thus sensitive to the prescribed search radius. A too small value might lead to a false splitting of the track, whereas a too large value would lead to more than one eddy within the searching area. As a first approximation, eddies are advected with the mean current. Considering a mean velocity of around 0.2 m/s (see e.g. Figure 5 in Wekerle et al. (2017)) and a daily mean model velocity field, a possible choice would be a search radius of 17 km. After performing sensitivity tests with different radii, we chose a radius of 14 km. This value reduced the number of occasions when several eddies were detected in the searching area. Furthermore, eddies with a lifetime shorter than 3 days were discarded.

For the eddy detection and tracking, we use daily model output for the time period 2006–2009 at 100 m depth. At this depth, the water mass lateral distribution is characterised by warm and salty AW in the eastern part of Fram Strait (in the WSC), and by



120 cold and fresh PW in its western part (in the EGC). Output from both models is interpolated to a regular grid (0.05° longitude
 x 0.01° latitude) which has approximately the same resolution as the original grids. Relative vorticity normalised by f at 100
 m depth on Jan 1st 2006 is shown in Figure 2, as well as eddies detected by the Nencioli et al. (2010) method overlaid on the
 simulated Okubo-Weiss parameter. Note that the colour only shows the area with $OW < -0.2\sigma_{OW}$, i.e. the area considered as
 vortices. In both models, the relative vorticity field exhibits strong eddy activity, particularly along the pathway of the main
 125 currents, WSC and EGC, along the Yermak and Svalbard branches and in the AW recirculation area. Apart from well defined
 eddies, the relative vorticity fields show lots of elongated filamentary structures reminiscent to what was found by von Appen
 et al. (2018). They seem to have a smaller scale in ROMS than in FESOM. Most of the eddies can be detected by both the
 Okubo-Weiss method and the Nencioli et al. (2010) method, while the Okubo-Weiss method detected a bit more eddies. Note
 that it was shown by Chaigneau et al. (2008) that the Okubo-Weiss criterion tends to over-detect eddies by comparison with
 130 eddies detected by experts.

2.4 Reynolds decomposition of eddy fluxes and and kinetic energy

To estimate the contributions of mesoscale eddy field to the flow variability, we decompose a variable x which can stand for
 velocity (u) or tracers (c) into a monthly mean (\bar{x}) and a daily-averaged fluctuating (x') component, $x = \bar{x} + x'$. We derive the
 time-mean eddy flux of the tracer c in the u velocity direction from the equality $\overline{c'u'} = \overline{c'u} - \bar{c}\bar{u}$. Similarly, time-averaged eddy
 135 kinetic energy (EKE) is computed as

$$EKE = \frac{1}{2}(\overline{u'^2} + \overline{v'^2}) = \frac{1}{2}(\overline{u^2} + \overline{v^2} - \bar{u}^2 - \bar{v}^2). \quad (2)$$

2.5 Energy budget

An energy budget can be obtained by expressing velocity as $\mathbf{u} = \bar{\mathbf{u}} + \mathbf{u}'$ as described in the previous section, inserting it in
 the momentum equation in the Boussinesq approximation, multiplying the equation with \mathbf{u}' , and time-averaging it. This leads
 to a conservation equation for EKE. The change of EKE in time is governed by the advection of eddies, energy transfer from
 140 mean kinetic energy (MKE) and eddy available potential energy (EPE) to EKE, and energy dissipation (vertical mixing and
 horizontal diffusion) (e.g. Olbers et al., 2012, chapter 12):

$$\frac{\partial \frac{1}{2}(\overline{u_1'^2} + \overline{u_2'^2})}{\partial t} + \underbrace{\frac{\partial \left(\frac{1}{2} \overline{u_j' u_i'^2} + \frac{1}{2} \overline{u_j' u_i'^2} + \frac{1}{\rho_0} \overline{u_j' p'} \right)}{\partial x_j}}_{\text{Transport}} = \underbrace{-\overline{u_j' u_i'} \frac{\partial u_i}{\partial x_j}}_{\text{MKE} \leftrightarrow \text{EKE}} + \underbrace{\overline{w' b'}}_{\text{EPE} \leftrightarrow \text{EKE}} + \underbrace{\overline{V_i' u_i'} + \overline{D_i' u_i'}}_{\text{Dissipation}}, \quad (3)$$

where $b = -\frac{g\rho}{\rho_0}$ is the buoyancy and D_i and V_i are horizontal and vertical dissipation terms. Cartesian tensor notation with
 summation convention has been used, with $i = 1, 2$ and $j = 1, 2, 3$. u_i is thus the horizontal component of the velocity vector
 u_j , and $u_3 = w$ is the vertical velocity. In this study, we diagnose the first two terms on the right hand side of the equation.
 145 They are the main source terms of EKE, and indicate barotropic and baroclinic instability.



3 Model assessment

For more than two decades, mooring measurements have been conducted across Fram Strait at around 79°N to monitor the exchange of water masses through this gateway (e.g. Beszczynska-Möller et al., 2012; von Appen et al., 2016; von Appen et al., 2019). To assess the overall model performance in reproducing the mean state and resolving the flow variability, we use
150 the observed hydrography as well as the velocity field and compare the latter in terms of power density spectra (PDS) and EKE to the model results.

The two models simulated very similar spatial distributions of water masses. The simulated mean temperature and salinity at 100 m depth reveal that the warm ($>5^{\circ}\text{C}$) and narrow WSC closely follows the 1000 m isobath along the Svalbard shelf break (Figure 3). Recirculation of AW mainly occurs north of the Boreas Basin (north of 78°N). The western part of Fram Strait
155 is characterised by cold and fresh polar outflow. The two models differ more significantly north of 80°N, with much warmer and saltier waters on the Yermak Plateau in FESOM. Compared to the mooring observations across Fram Strait, both models correctly represent the main water masses. ROMS shows a slightly cold bias which is not present in FESOM (ROMS: root mean square (rms) error of 1.28°C , FESOM: rms error of 0.49°C), and has earlier been identified to be associated with a cold bias in the A4 model that provides the inflow boundary conditions for S800 (Hattermann et al., 2016). The simulated water
160 mass in FESOM, particularly in central and eastern Fram Strait, is slightly too saline, whereas it is slightly too fresh in ROMS. The overall rms error in salinity is 0.26 and 0.31 in ROMS and FESOM, respectively.

For the comparison of velocity time series, current meter data from three moorings located in the WSC and three moorings located in the EGC (for locations see Figure 1) deployed during the time period 2006–2009 were used (von Appen et al., 2019). Daily averages of measured velocity at 75 m depth were calculated. Note that there are slight variations in the depth between
165 the individual deployment years. The observed mean speed averaged over WSC and EGC moorings at 75 m depth is 0.22 m/s and 0.13 m/s respectively, while the mean speed of ROMS/FESOM at the mooring locations is 0.24/0.20 m/s and 0.16/0.12 m/s respectively. Power density spectra of the speed time series from the observations (linearly interpolated smaller gaps, e.g. between individual deployments) and from the models were estimated via the Thomson multitaper method (Figure 4). For the estimated PDS, the median in $\log_{10}(0.05/\text{day})$ frequency steps was calculated for frequencies between $-1.2/\text{day}$ and $-0.35/\text{day}$,
170 and the slopes were fitted to those binned values. The slopes of the observations are ~ -1.7 and ~ -1.5 for WSC and EGC moorings, respectively, while ROMS/FESOM showed slopes of $\sim -1.7/\sim -2.1$ and $\sim -2.2/\sim -2.7$ respectively. The difference between the models is larger at high frequency, which might be related to the fact that tides were simulated in ROMS.

A seasonal cycle of EKE in 75 m depth computed from current meter data of moorings deployed across Fram Strait is shown in Figure 5a. The highest level of EKE is reached in the winter months (January–March), and lowest values are reached in
175 early autumn (September–November). There is a strong lateral gradient from west to east, with a much higher level of EKE in the eastern part of Fram Strait, the WSC region. Both models well reproduce the observed seasonal and spatial variations of EKE (Figure 5b and c), except that the observation shows a higher EKE level in the central Fram Strait than the models.



4 Eddy properties

4.1 Eddy spatial distribution and polarisation

180 During the time period 2006–2009, altogether 218,213 eddies were detected in the area 8°W – 20°E / 76°N – 82°N in ROMS (and thus 149 eddies per day), with slightly more cyclones (54%) than anticyclones. The result is very similar in FESOM, with 55% of the 244,811 detected eddies (168 per day) being cyclones. The tracking algorithm then revealed that these eddies belong to 30539 and 39040 tracks for ROMS and FESOM, respectively. In both simulations, the eddy density is highest in the eastern and central part of Fram Strait (Figure 6a,b). In contrast, the eddy density is low in the western part of Fram Strait and on the
185 East Greenland continental shelf, that is, in areas covered by sea ice year-round. Comparing FESOM and ROMS, there are fewer eddies detected in that region in FESOM, which is also reflected in lower EKE values in the western part of Fram Strait than in ROMS (Figure 5). Both models show a consistent pattern in the distribution of cyclones vs. anticyclones, which has strong regional differences (Figure 6c and d). Over the Svalbard shelf and along the East Greenland continental shelf break, cyclones are predominant. Anticyclones dominate along the main pathway of the WSC (along the 1000 m isobath), over the
190 Yermak Plateau, and along the Svalbard branch.

4.2 Eddy size

In this study we compute the eddy radius as average distance from the eddy centre to the eddy boundary, which is defined by the outermost closed contour of the stream function field. Eddies detected in both models are relatively small, with 95%/92% of cyclones and 92%/87% of anticyclones in ROMS/FESOM having a radius below 10 km (Figure 7a). The mean/median radius
195 for ROMS and FESOM is 4.9/4.1 and 5.6/4.7 km, respectively (Table 2). Eddies simulated in FESOM are thus slightly larger than in ROMS. The eddy radius compares well with the Rossby radius of deformation (~ 4 – 6 km in summer and smaller values in winter (von Appen et al., 2016)). This suggests that baroclinic instability is likely the main mechanism of eddy generation, which will be further investigated in Section 5. In both simulations, cyclones are slightly smaller than anticyclones (Table 2).

4.3 Eddy intensity

200 The mean/median intensity of eddies (expressed as the absolute value of relative vorticity normalised by f , averaged over all detected eddies) simulated by ROMS and FESOM is 0.4/0.36 and 0.28/0.24, respectively. Eddies simulated by FESOM are thus weaker than eddies simulated by ROMS (see also Figure 7b and Table 2). The proportion of eddies with absolute values below 0.3 is larger for FESOM (63%) than for ROMS (38%). Cyclones are slightly more intensive and have a larger standard deviation than anticyclones in both models (Table 2).

205 4.4 Eddy lifetime and travel distance

The duration over which eddies are continuously detected by the employed method is on average 10 and 11 days in ROMS and FESOM, respectively (Figure 7d). 85%/82% of eddies detected in ROMS/FESOM have lifetimes below 15 days, whereas



only 4%/6% of eddies detected in ROMS/FESOM have lifetimes above 30 days. Pathways of these long-living eddies will be analysed in the next section. Note that the eddy lifetime may be longer if one considers that eddies likely can exist for some time before and after being detected as an eddy by the tracking method. Also, a false splitting of the track could occur if the eddy moved relatively fast in combination with a too small searching area. In both simulations, there is no significant difference in lifetime regarding polarisation. They are very similar regarding travel distance. On average, eddies travel around 34 and 35 km in ROMS and FESOM, respectively (Table 2). Again, there is no significant difference in travel distance regarding polarisation (Figure 7e). Compared to eddies generated e.g. in the Gulf Stream region, the lifetime of Fram Strait eddies is rather short (Kang and Curchitser, 2013).

4.5 Eddy pathways

Eddy pathways are investigated by focusing only on long-living eddies, e.g. eddies with lifetime of more than 30 days, and by classifying them by generation areas (Figures 8 and 9). In both simulations, eddies generated on the Svalbard shelf have very distinct travel pathways for cyclones and anticyclones, which is consistent with their distribution (Figure 6e and f). Cyclones tend to stay on the shelf, and populate the narrow Svalbard fjords. Anticyclones in contrast leave the shallow shelf area and tend to travel westward into the deep basin. As shown in Figure 7c, more cyclones (31% and 25% in ROMS and FESOM, respectively) are detected in shallow areas with water depths less than 500 m than anticyclones (21% and 19% in ROMS and FESOM, respectively). Note that as the number of detected eddies on the East Greenland shelf is relatively small in both simulations, most eddies detected in shallow areas are located on the Svalbard shelf.

Anticyclones generated in the WSC core region, here defined approximately as the area between the 500 m and 2000 m isobaths, show longer travel pathways than cyclones. In both simulations, most of them travel westward along the recirculation pathway north of the Molloy Deep (Hattermann et al., 2016), and some even continue southward along the East Greenland continental shelf break. Some eddies travel northward along the western rim of the Yermak plateau or recirculate around the Molly Deep, while only few trajectories deviate westward south of 79°N in both models.

The asymmetric pathways of eddies generated on the Svalbard shelf and in the WSC core region can have dynamical reasons. As described by Cushman-Roisin (1994, Chapter 17), fluid parcels surrounding a rotating eddy are stretched when they move to deeper waters and thus acquire relative vorticity. In contrast, when moving to shallower waters, on the flank of the eddy the surrounding fluid is squeezed and thus relative vorticity is decreased. This results in a secondary drift of the vortices, with cyclones moving towards shallower regions and anticyclones moving to deeper regions. Morrow et al. (2004), based on satellite altimetry, showed that this dynamical reasoning can explain the diverging pathways of cyclones and anticyclones in different ocean basins.

Tracks of long-living eddies generated in the southern central Fram Strait, in particular those simulated in ROMS, show a high density of anticyclones in the Boreas Basin, the region between 0°EW–5°E, 76°N–77°N. More anticyclones appear to be trapped in this depression, a similar situation as occurring in the Lofoten Basin (Raj et al., 2016; Volkov et al., 2015). As in the case of eddies generated along the Svalbard shelf break, the clustering of anticyclones can be explained by the dynamical reason described above (anticyclones move towards the deeper basin, thus the centre of a depression). Eddies generated in northern



central Fram Strait tend to travel westward, then follow the East Greenland continental shelf break. Particularly, anticyclones travel westward between the northern rim of the Boreas Basin and the Molloy Deep, contributing to the AW recirculation.

Regarding eddies present in northern Fram Strait, both ROMS and FESOM show a high density along the western flank of the Yermak Plateau. Additionally, ROMS shows more long-living (>30 days lifetime) eddies west of the Plateau (Figure 6a-d) than FESOM (Figures 9 and 8). Eddies in this region have earlier been identified to occur with a different seasonality than would be expected from changes in baroclinic instability of the boundary current that explains the seasonality in eddy occurrence along other parts of the shelf break (Crews et al., 2019). A difference between the two models is the inclusion of tidal forcing in ROMS. The circulation and water mass transformations above the Yermak plateau are known to be strongly influenced by barotropic to baroclinic tidal conversion and mixing at the semi-diurnal critical latitude (Fer et al., 2015), that may also explain the enhanced eddy generation in this region in ROMS. As revealed by FESOM, more cyclones tend to follow the Svalbard Branch, whereas more anticyclones tend to follow the Yermak Branch.

4.6 Vertical extent and hydrographic properties

We determined the vertical extent of eddies detected in 100 m depth with lifetime above 30 days by calculating relative vorticity/ f in the eddy centres in the water column (Figure 10). In addition, temperature and salinity anomalies were calculated in the same way to study the hydrographic properties of eddies, with anomalies computed relative to the mean value for the month. This was done for eddies generated in the five different regions shown in Figure 1. Profiles of relative vorticity are relatively similar in ROMS and FESOM, with most negative/positive (i.e. strongest vortices) values for anticyclones/cyclones generated in the WSC region and central Fram Strait.

The hydrographic conditions in regions WSC, central southern Fram Strait and Yermak/Svalbard Branch are characterised by warm and salty AW (Figure 3). These regions are temperature-stratified and unstable in salinity. Anticyclones generated there carry anomalously warm and salty and thus lighter waters and have depressed isopycnals, whereas cyclones carry anomalously cold and fresh and thus denser waters and have raised isopycnals (Figure 10). The western Fram Strait is characterised by cold and fresh PW, and is salinity stratified. The transition from a temperature stratified to a salinity stratified regime in the different regions may partly explain the difference in properties between ROMS and FESOM.

5 Energetics in the eastern Fram Strait

We now analyse the source of EKE as simulated in ROMS and FESOM. We focus here on the eastern side of Fram Strait, which is the most energetic region (Figure 5). As described in section 2.5, the change of EKE in time is governed by the advection of eddies, energy transfer from mean kinetic energy (MKE) and eddy available potential energy (EPE) to EKE, and energy dissipation. In this study we analyse only the first two terms on the right hand side of the EKE conservation equation (Eq. 3), which are the main source terms for EKE and are related to barotropic and baroclinic instability.



5.1 Barotropic instability

The transfer of MKE to EKE is related to barotropic instability. It can be expressed as the sum of two terms, the product of horizontal eddy Reynolds stress and horizontal mean shear, and the product of vertical eddy Reynolds stress and vertical mean shear. Strong velocity shear thus support barotropic instability. Here we consider only terms that contain horizontal derivatives, and assume that the terms with vertical derivatives play a minor role (as shown for the Gulf Stream region by Gula et al. (2015)). In the two models, the energy conversion between MKE and EKE is directed in both ways: it shows an alternating pattern, with positive values indicating conversion from MKE to EKE and negative values indicating conversion from EKE to MKE (Figure 11a,b)¹. The alternating pattern is very similar between the both models, with consistent locations and magnitude of positive and negative energy transfer. The energy transfer occurs mainly along the pathway of the WSC core, which is located approximately along the 500–1000 m isobaths (see also Figure 3).

The relatively similar pattern in both models suggests that there is a strong influence of bathymetry, which determines positive and negative spots of energy conversion. A necessary condition for barotropic instability is that $\beta - \partial_{yy}\bar{u}$ vanishes within the domain, where $u = \bar{u}(y)$ is a zonal current with arbitrary meridional profile (e.g. Cushman-Roisin, 1994). The planetary potential vorticity is weak and can be ignored in polar regions, so we only consider the topographic β , with $\beta = -\frac{f}{H}\nabla H$ quantifying the change in potential vorticity across the bathymetry and H and ∇H being the water depth and its horizontal gradient. A map of the topographic β west of Svalbard reveals large values along the Svalbard shelf break (Figure 12a). We take the monthly mean meridional velocity v from FESOM as an approximation of the along-stream velocity, and compute $\beta - \partial_{xx}v$ (Figure 12b). In many places along the Svalbard shelf break, β is much larger than $\partial_{xx}v$. However, in some places, e.g. at the entrance of Kongsfjorden (79°N) and Isfjorden (78°10'N) and along the 250 m isobath at around 80°N, $\beta - \partial_{xx}v$ changes sign. These regions are characterised by positive values of energy conversion in both models, indicating active barotropic instability there. This is comparable to the Norwegian continental slope off the Lofoten islands, where barotropic instability becomes particularly important in regions with steep bottom slope (Ghaffari et al., 2018).

5.2 Baroclinic instability

For baroclinic instability to be active, a horizontal density gradient must be present to provide available potential energy which can be converted to EKE. This transfer from EPE to EKE can be expressed as the mean vertical eddy buoyancy flux (Eq. 3). In contrast to barotropic instability, the energy conversion between EPE and EKE in FESOM and ROMS is directed mostly one way, with mainly positive values revealing conversion from EPE to EKE (Figure 11c,d). As the eastern Fram Strait is temperature-stratified, it is mainly the vertical eddy temperature flux that contributes to vertical eddy buoyancy flux (Hattermann et al., 2016, their Figure 3d). In both models, baroclinic instability is strongest between the 1000 m and 2000 m isobaths in eastern Fram Strait. The values are slightly weaker in FESOM than in ROMS. The weaker baroclinic instability in FESOM is also reflected by the fact that detected eddies are characterised by lower values of absolute relative vorticity (Figure

¹Note that there has been an error in the computation of the MKE to EKE conversion term shown in Figure 14a of Wekerle et al. (2017). Figure 11b shows the correct pattern.



7b). Between the 500 m and 1000 m isobaths, both models show patches of negative vertical eddy buoyancy fluxes. Usually, those patches indicate regions where eddy fluxes interact with the sloping topography to lift dense water onto the continental shelf (Tverberg and Nøst, 2009), with upward sloping isopycnals near the seafloor that locally enhance the APE of the mean field.

A necessary condition for baroclinic instability is that the cross-stream gradient of Ertel potential vorticity (PV) changes sign with depth (e.g. Spall and Pedlosky, 2008). Ertel PV Π is defined as

$$\begin{aligned} \Pi &= (f\mathbf{k} + \nabla \times \mathbf{u})\nabla b \\ 310 \quad &= \underbrace{f \partial_z b}_{\text{vertical stretching}} + \underbrace{(\partial_y w - \partial_z v) \partial_x b + (\partial_x w - \partial_z u) \partial_y b}_{\text{tilting vorticity}} + \underbrace{(\partial_x v - \partial_y u) \partial_z b}_{\text{relative vorticity}}. \end{aligned}$$

Here we compute Π from simulated long-term mean velocity $\mathbf{u} = (u, v, w)$ and buoyancy b , and neglect the small terms containing derivatives of vertical velocity w . Figure 13 shows the Ertel PV and its gradient in zonal direction for two sections across the Svalbard shelf break (78°N and 78°50'N) for the FESOM simulation. The dominant term is the vertical stretching term, with a smaller contribution from the relative vorticity terms. The tilting terms are one order of magnitude smaller (Figure not shown). At both sections, the cross-stream gradient reveals a change in sign with depth, indicating that the mean current is baroclinically unstable. This is in agreement with studies by Teigen et al. (2011) and von Appen et al. (2016), and our simulated energy conversions (Figure 11c,d).

6 Discussion

Despite their very fine resolution, ROMS and FESOM simulate a weaker variability in velocity than the observed in terms of the power density spectrum (Figure 4). This might indicate that the resolution used is still insufficient to well resolve all the mesoscale eddies in the presence of numerical dissipation. Part of the variability revealed by the power density spectrum can also be attributed to the atmospheric forcing. Although the forcing datasets are different in the two cases, both of them are derived from relatively coarse reanalysis products (in particular, COREv.2 used in the FESOM simulation has a zonal resolution of approximately 1.875°) and may miss part of small-scale variability. A topic for further research is to clarify the importance of these factors.

Snapshots of simulated relative vorticity (Figure 2) and the baroclinic energy transfer (Figure 11c,d) suggest that the model effective resolution (Soufflet et al., 2016) in FESOM might be slightly lower than in ROMS. First, the grid size is slightly larger for FESOM (1 km vs 800 m for ROMS). This small difference in the grid size (20%) might matter as both numerical dissipation and explicit viscosity decrease with the grid size. In both models, biharmonic viscosity which scales with grid size cubed is applied. Second, FESOM1.4 is based on a collocated discretisation (an analog of Arakawa A-grid), whereas a staggered Arakawa C-grid is employed by ROMS. Because of pressure gradient averaging required by collocated discretisations, the effective resolution could be reduced. The collocated discretisation of FESOM also requires to use the no-slip boundary condition, which implies more dissipation along the boundary as well. Third, FESOM relies on implicit time stepping for external mode whereas ROMS uses a specially selected split-explicit method (see, e.g. Soufflet et al. (2016)) which is less



335 dissipative. There might be other reasons for the difference in the simulated EKE in certain regions between the two models.
In particular, a higher EKE level in western Fram Strait in ROMS might be related to the difference in the simulated sea ice.
Sea ice could damp eddies through the ocean-ice stress.

In addition to the high similarity in the eddy properties, both models exhibit a very similar pattern in barotropic energy
conversion in eastern Fram Strait (Figure 11a,b). The degree of similarity is quite surprising, given that FESOM uses z-levels
340 in the vertical whereas ROMS relies on terrain-following coordinates, which might lead to differences in topographic steering.
Topography is bi-linearly interpolated to grid points and only smoothed over the 2d-stencil of nearest vertices in FESOM. In
contrast, ROMS requires a smoother bathymetry.

Our instability analysis provides information that can be used to develop parameterisations for coarse resolution climate
models. A commonly used eddy parameterisation is the one developed by Gent and McWilliams (1990), which mixes tracers
345 along local potential density, and thus flatten isopycnals. However, our analysis shows that there are areas with conversion
from EKE to MKE (Figure 11a,b) associated with a steepening of isopycnals, which is not taken into account by the Gent-
McWilliams parameterisation.

7 Conclusions

Several studies (e.g. Hattermann et al., 2016; Wekerle et al., 2017) indicate that mesoscale eddies are an important ingredient
350 of ocean dynamics in the Fram Strait, determining the recirculation of AW and also the amount of oceanic heat that enters the
Arctic Ocean. However, the eddy properties have not been thoroughly investigated before.

Based on the results of two eddy-resolving ocean-sea ice models, ROMS and FESOM, we examined the properties and
generation mechanisms of mesoscale eddies in Fram Strait. We found that the models agree with each other with respect to the
modelled circulation, hydrography and eddy characteristics. They simulate rather short-living eddies (lifetime is on average
355 10–11 days), with a very slight dominance of cyclones (ROMS: 54%, FESOM: 55%). Cyclones and anticyclones show very
distinct travel pathways, e.g., cyclones generated on the shallow Svalbard shelf tend to stay there, whereas anticyclones tend to
travel offshore into the deep basin. More anticyclones tend to be trapped in deep depressions. Mean eddy radius is 5.0–6.0 km,
which compares well with the first baroclinic Rossby radius of deformation in this region. On average, eddies travel around 35
km in both models. Eddy cores are located at about 100 m depth on average. Cyclones are predominantly cold eddies, while
360 anticyclones are predominantly warm eddies.

The models also agree on mechanisms driving eddy generation, with consistent patterns of conversions to EKE from the
mean kinetic and eddy available potential energies. The small size of eddies explains why a very high (1 km or finer) resolution
is needed to simulate them. The good agreement on eddy generation and properties despite the very different numerics of
FESOM (unstructured horizontal grid with vertical z-levels) and ROMS (regular horizontal grid with a terrain following vertical
365 coordinate) gives us confidence in their ability to realistically simulate eddy processes. The similarities of the simulated eddy
fields despite the different sea ice components, surface and boundary conditions of the two models further imply that the
observed eddy characteristics are likely a result of the fundamental dynamics associated with the ocean mean state in this



region, rather than reflecting the variability imposed by an arbitrary external forcing. Some differences between the two models are also identified in this work, including the intensity of eddies and the rates of energy conversion, which require more
370 dedicated research to better understand the reasons.



Author contributions. CW, QW and SD contributed the FESOM data and analysis, TH and LC contributed the ROMS data, and WJvA contributed the observational data. All authors discussed the content of the manuscript and contributed to the interpretation of data and writing of the manuscript.

Competing interests. We declare that no competing interests are present.

375 *Acknowledgements.* This work was supported by the AWI FRAM (FRontiers in Arctic marine Monitoring) program (CW and WJvA). TH
acknowledges financial support from the Norwegian Research Council project 280727. LC acknowledges support from the Fram Centre
"Arctic Ocean" flagship project "Mesoscale modeling of Ice, Ocean, and Ecology of the Arctic Ocean (ModOIE)" and Office of Naval
Research grant number N00014-18-1-2694. This material is based upon work supported by the National Science Foundation Graduate
Research Fellowship Program under Grant No. DGE-1762114. Any opinions, findings, and conclusions or recommendations expressed in
380 this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation. The FESOM simulation
was performed at the North-German Supercomputing Alliance (HLRN). The ROMS simulation was performed under the NOTUR hpc project
mn9238k.



References

- Beszczynska-Möller, A., Fahrbach, E., Schauer, U., and Hansen, E.: Variability in Atlantic water temperature and transport at the entrance to the Arctic Ocean, 1997–2010, *ICES J. Mar. Sci.*, 69, 852–863, <https://doi.org/10.1093/icesjms/fss056>, 2012.
- Budgell, W. P.: Numerical simulation of ice-ocean variability in the Barents Sea region, *Ocean Dynamics*, 55, 370–387, <https://doi.org/10.1007/s10236-005-0008-3>, 2005.
- Chaigneau, A., Gizolme, A., and Grados, C.: Mesoscale eddies off Peru in altimeter records: Identification algorithms and eddy spatio-temporal patterns, *Progress in Oceanography*, 79, 106 – 119, <https://doi.org/10.1016/j.pocean.2008.10.013>, 2008.
- 385 Cherian, D. A. and Brink, K. H.: Shelf Flows Forced by Deep-Ocean Anticyclonic Eddies at the Shelf Break, *Journal of Physical Oceanography*, 48, 1117–1138, <https://doi.org/10.1175/JPO-D-17-0237.1>, 2018.
- Crews, L., Sundfjord, A., Albretsen, J., and Hattermann, T.: Mesoscale Eddy Activity and Transport in the Atlantic Water Inflow Region North of Svalbard, *Journal of Geophysical Research: Oceans*, 123, 201–215, <https://doi.org/10.1002/2017JC013198>, 2018.
- Crews, L., Sundfjord, A., and Hattermann, T.: How the Yermak Pass Branch Regulates Atlantic Water Inflow to the Arctic Ocean, *Journal of Geophysical Research: Oceans*, 124, 267–280, <https://doi.org/10.1029/2018JC014476>, 2019.
- 395 Cushman-Roisin, B.: *Introduction to Geophysical Fluid Dynamics*, Prentice Hall, 1994.
- Dai, A., Qian, T., Trenberth, K. E., and Milliman, J. D.: Changes in Continental Freshwater Discharge from 1948 to 2004, *J. Climate*, 22, 2773–2792, <https://doi.org/10.1175/2008JCLI2592.1>, 2009.
- Danilov, S., Wang, Q., Timmermann, R., Iakovlev, N., Sidorenko, D., Kimmritz, M., Jung, T., and Schröter, J.: Finite-Element Sea Ice Model (FESIM), version 2, *Geosci. Model Dev.*, 8, 1747–1761, <https://doi.org/10.5194/gmd-8-1747-2015>, 2015.
- 400 de Steur, L., Hansen, E., Gerdes, R., Karcher, M., Fahrbach, E., and Holfort, J.: Freshwater fluxes in the East Greenland Current: A decade of observations, *Geophys. Res. Lett.*, 36, <https://doi.org/10.1029/2009GL041278>, 2009.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quarterly Journal of the Royal Meteorological Society*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- 405 Egbert, G. D. and Erofeeva, S. Y.: Efficient Inverse Modeling of Barotropic Ocean Tides, *Journal of Atmospheric and Oceanic Technology*, 19, 183–204, [https://doi.org/10.1175/1520-0426\(2002\)019<0183:EIMOBO>2.0.CO;2](https://doi.org/10.1175/1520-0426(2002)019<0183:EIMOBO>2.0.CO;2), 2002.
- Eldevik, T., Nilsen, J. E. Ø., Iovino, D., Olsson, K. A., Sandø, A. B., and Drange, H.: Observed sources and variability of Nordic Seas overflow, *Nature Geoscience*, 2, 406–410, <https://doi.org/10.1038/NGEO518>, 2009.
- Fer, I., Müller, M., and Peterson, A. K.: Tidal forcing, energetics, and mixing near the Yermak Plateau, *Ocean Science*, 11, 287–304, <https://doi.org/10.5194/os-11-287-2015>, 2015.
- 415 Gent, P. and McWilliams, J.: Isopycnal Mixing in Ocean Circulation Models, *J. Phys. Oceanogr.*, 20, 150–155, 1990.
- Ghaffari, P., Isachsen, P. E., Nøst, O. A., and Weber, J. E.: The Influence of Topography on the Stability of the Norwegian Atlantic Current off Northern Norway, *Journal of Physical Oceanography*, 48, 2761–2777, <https://doi.org/10.1175/JPO-D-17-0235.1>, 2018.
- Gula, J., Molemaker, M. J., and McWilliams, J. C.: Gulf Stream Dynamics along the Southeastern U.S. Seaboard, *J. Phys. Oceanogr.*, 45, 690–715, <https://doi.org/10.1175/JPO-D-14-0154.1>, 2015.



- 420 Haidvogel, D., Arango, H., Budgell, W., Cornuelle, B., Curchitser, E., Lorenzo, E. D., Fennel, K., Geyer, W., Hermann, A., Lanerolle, L., Levin, J., McWilliams, J., Miller, A., Moore, A., Powell, T., Shchepetkin, A., Sherwood, C., Signell, R., Warner, J., and Wilkin, J.: Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the Regional Ocean Modeling System, *Journal of Computational Physics*, 227, 3595 – 3624, <https://doi.org/10.1016/j.jcp.2007.06.016>, 2008.
- Hattermann, T., Isachsen, P. E., von Appen, W.-J., Albretsen, J., and Sundfjord, A.: Eddy-driven recirculation of Atlantic Water in Fram Strait, *Geophys. Res. Lett.*, 43, <https://doi.org/10.1002/2016GL068323>, 2016.
- 425 Isern-Fontanet, J., Garcia-Ladona, E., and Font, J.: Vortices of the Mediterranean Sea: An Altimetric Perspective, *Journal of Physical Oceanography*, 36, 87–103, <https://doi.org/10.1175/JPO2826.1>, 2006.
- Johannessen, J. A., Johannessen, O. M., Svendsen, E., Shuchman, R., Manley, T., Campbell, W. J., Josberger, E. G., Sandven, S., Gascard, J. C., Olaussen, T., Davidson, K., and Van Leer, J.: Mesoscale eddies in the Fram Strait marginal ice zone during the 1983 and 1984 Marginal Ice Zone Experiments, *J. Geophys. Res. Oceans*, 92, 6754–6772, <https://doi.org/10.1029/JC092iC07p06754>, 1987a.
- 430 Johannessen, O. M., Johannessen, J. A., Svendsen, E., Shuchman, R. A., Campbell, W. J., and Josberger, E.: Ice-Edge Eddies in the Fram Strait Marginal Ice Zone, *Science*, 236, 427–429, <https://doi.org/10.1126/science.236.4800.427>, 1987b.
- Kang, D. and Curchitser, E. N.: Gulf Stream eddy characteristics in a high-resolution ocean model, *Journal of Geophysical Research: Oceans*, 118, 4474–4487, <https://doi.org/10.1002/jgrc.20318>, 2013.
- 435 Kawasaki, T. and Hasumi, H.: The inflow of Atlantic water at the Fram Strait and its interannual variability, *J. Geophys. Res. Oceans*, 121, 502–519, <https://doi.org/10.1002/2015JC011375>, 2016.
- Large, W. and Yeager, S.: The global climatology of an interannually varying air-sea flux data set, *Clim. Dynam.*, 33, 341–364, <https://doi.org/10.1007/s00382-008-0441-3>, 2008.
- Mahadevan, A.: The Impact of Submesoscale Physics on Primary Productivity of Plankton, *Annual Review of Marine Science*, 8, 161–184, <https://doi.org/10.1146/annurev-marine-010814-015912>, 2016.
- 440 Manucharyan, G. E. and Thompson, A. F.: Submesoscale Sea Ice-Ocean Interactions in Marginal Ice Zones, *Journal of Geophysical Research: Oceans*, 122, 9455–9475, <https://doi.org/10.1002/2017JC012895>, 2017.
- Morrow, R., Birol, F., Griffin, D., and Sudre, J.: Divergent pathways of cyclonic and anti-cyclonic ocean eddies, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL020974>, 2004.
- 445 Nencioli, F., Dong, C., Dickey, T., Washburn, L., and McWilliams, J. C.: A Vector Geometry-Based Eddy Detection Algorithm and Its Application to a High-Resolution Numerical Model Product and High-Frequency Radar Surface Velocities in the Southern California Bight, *Journal of Atmospheric and Oceanic Technology*, 27, 564–579, <https://doi.org/10.1175/2009JTECHO725.1>, 2010.
- Okubo, A.: Horizontal dispersion of floatable particles in the vicinity of velocity singularities such as convergences, *Deep Sea Research and Oceanographic Abstracts*, 17, 445 – 454, [https://doi.org/10.1016/0011-7471\(70\)90059-8](https://doi.org/10.1016/0011-7471(70)90059-8), 1970.
- 450 Olbers, D., Willebrand, J., and Eden, C.: *Ocean Dynamics*, Springer-Verlag Berlin Heidelberg, <https://doi.org/10.1007/978-3-642-23450-7>, 2012.
- Polyakov, I. V., Pnyushkov, A. V., and Timokhov, L.: Warming of the Intermediate Atlantic Water of the Arctic Ocean in the 2000s, *J. Climate*, 25, 8362–8370, <https://doi.org/10.1175/JCLI-D-12-00266.1>, 2012.
- Raj, R. P., Johannessen, J. A., Eldevik, T., Nilsen, J. E. A., and Halo, I.: Quantifying mesoscale eddies in the Lofoten Basin, *Journal of Geophysical Research: Oceans*, 121, 4503–4521, <https://doi.org/10.1002/2016JC011637>, 2016.
- 455 Rudels, B.: The thermohaline circulation of the Arctic Ocean and the Greenland Sea, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 352, 287–299, <https://doi.org/10.1098/rsta.1995.0071>, 1995.



- Schaffer, J., von Appen, W.-J., Dodd, P. A., Hofstede, C., Mayer, C., de Steur, L., and Kanzow, T.: Warm water pathways toward Nioghalvfjerdingsfjorden Glacier, Northeast Greenland, *Journal of Geophysical Research: Oceans*, 122, 4004–4020, <https://doi.org/10.1002/2016JC012462>, 2017.
- Schauer, U., Fahrbach, E., Østerhus, S., and Rohardt, G.: Arctic warming through the Fram Strait: Oceanic heat transport from 3 years of measurements, *J. Geophys. Res.*, 109, 2004.
- Shchepetkin, A. F. and McWilliams, J. C.: The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Modelling*, 9, 347 – 404, <https://doi.org/10.1016/j.ocemod.2004.08.002>, 2005.
- 465 Shchepetkin, A. F. and McWilliams, J. C.: Correction and commentary for "Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the regional ocean modeling system" by Haidvogel et al., *J. Comp. Phys.* 227, pp. 3595–3624, *Journal of Computational Physics*, 228, 8985 – 9000, <https://doi.org/10.1016/j.jcp.2009.09.002>, 2009.
- Smith, D. C., Morison, J. H., Johannessen, J. A., and Untersteiner, N.: Topographic generation of an eddy at the edge of the East Greenland Current, *J. Geophys. Res. Oceans*, 89, 8205–8208, <https://doi.org/10.1029/JC089iC05p08205>, 1984.
- 470 Soufflet, Y., Marchesiello, P., Lemarié, F., Jouanno, J., Capet, X., Debreu, L., and Benshila, R.: On effective resolution in ocean models, *Ocean Modelling*, 98, 36 – 50, <https://doi.org/10.1016/j.ocemod.2015.12.004>, 2016.
- Spall, M. A. and Pedlosky, J.: Lateral Coupling in Baroclinically Unstable Flows, *Journal of Physical Oceanography*, 38, 1267–1277, <https://doi.org/10.1175/2007JPO3906.1>, 2008.
- Storkey, D., Blockley, E. W., Furner, R., Guiavarc'h, C., Lea, D., Martin, M. J., Barciela, R. M., Hines, A., Hyder, P., and Sid-
475 dorn, J. R.: Forecasting the ocean state using NEMO: The new FOAM system, *Journal of Operational Oceanography*, 3, 3–15, <https://doi.org/10.1080/1755876X.2010.11020109>, 2010.
- Sundfjord, A., Albretsen, J., Kasajima, Y., Skogseth, R., Kohler, J., Nuth, C., Skaróhamar, J., Cottier, F., Nilsen, F., Asplin, L., Gerland, S., and Torsvik, T.: Effects of glacier runoff and wind on surface layer dynamics and Atlantic Water exchange in Kongsfjorden, Svalbard; a model study, *Estuarine, Coastal and Shelf Science*, 187, 260 – 272, <https://doi.org/10.1016/j.ecss.2017.01.015>, 2017.
- 480 Teigen, S. H., Nilsen, F., Skogseth, R., Gjevik, B., and Beszczynska-Möller, A.: Baroclinic instability in the West Spitsbergen Current, *J. Geophys. Res. Oceans*, 116, <https://doi.org/10.1029/2011JC006974>, 2011.
- Tverberg, V. and Nøst, O. A.: Eddy overturning across a shelf edge front: Kongsfjorden, west Spitsbergen, *Journal of Geophysical Research: Oceans*, 114, <https://doi.org/10.1029/2008JC005106>, 2009.
- Volkov, D. L., Kubryakov, A. A., and Lumpkin, R.: Formation and variability of the Lofoten basin vortex in a high-resolution ocean model, *Deep Sea Research Part I*, 105, 142–157, <https://doi.org/10.1016/j.dsr.2015.09.001>, 2015.
- 485 von Appen, W.-J., Schauer, U., Hattermann, T., and Beszczynska-Möller, A.: Seasonal Cycle of Mesoscale Instability of the West Spitsbergen Current, *J. Phys. Oceanogr.*, 46, 1231–1254, <https://doi.org/10.1175/JPO-D-15-0184.1>, 2016.
- von Appen, W.-J., Wekerle, C., Hehemann, L., Schourup-Kristensen, V., Konrad, C., and Iversen, M. H.: Observations of a Submesoscale Cyclonic Filament in the Marginal Ice Zone, *Geophysical Research Letters*, 45, 6141–6149, <https://doi.org/10.1029/2018GL077897>, 2018.
- 490 von Appen, W.-J., Beszczynska-Möller, A., Schauer, U., and Fahrbach, E.: Physical oceanography and current meter data from moorings F1-F14 and F15/F16 in the Fram Strait, 1997-2016, <https://doi.org/10.1594/PANGAEA.900883>, 2019.
- Wang, Q., Danilov, S., and Schröter, J.: Finite element ocean circulation model based on triangular prismatic elements, with application in studying the effect of topography representation, *J. Geophys. Res.*, 113, 2008.



- 495 Wang, Q., Danilov, S., Sidorenko, D., Timmermann, R., Wekerle, C., Wang, X., Jung, T., and Schröter, J.: The Finite Element Sea Ice-Ocean Model (FESOM) v.1.4: formulation of an ocean general circulation model, *Geosci. Model Dev.*, 7, 663–693, <https://doi.org/10.5194/gmd-7-663-2014>, 2014.
- Weiss, J.: The dynamics of enstrophy transfer in two-dimensional hydrodynamics, *Physica D: Nonlinear Phenomena*, 48, 273 – 294, [https://doi.org/10.1016/0167-2789\(91\)90088-Q](https://doi.org/10.1016/0167-2789(91)90088-Q), 1991.
- 500 Wekerle, C., Wang, Q., von Appen, W.-J., Danilov, S., Schourup-Kristensen, V., and Jung, T.: Eddy-Resolving Simulation of the Atlantic Water Circulation in the Fram Strait With Focus on the Seasonal Cycle, *J. Geophys. Res. Oceans*, <https://doi.org/10.1002/2017JC012974>, 2017.
- Wilson, N., Straneo, F., and Heimbach, P.: Satellite-derived submarine melt rates and mass balance (2011–2015) for Greenland’s largest remaining ice tongues, *The Cryosphere*, 11, 2773–2782, <https://doi.org/10.5194/tc-11-2773-2017>, 2017.



Table 1. Characteristics of the FESOM and ROMS configurations used in this study.

	FESOM	ROMS
Numerical method	finite elements	finite differences
Horizontal grid	A-grid (P1-P1 scheme)	C-grid
Vertical coordinate	z-levels	terrain following levels
Domain	global	regional
Horizontal mixing scheme	biharmonic Smagorinsky	biharmonic
Vertical mixing scheme	KPP	KPP
Tides	no	yes

Table 2. Mean properties and their standard deviation (in brackets) for all eddies generated in the area 8°W–20°E/76°N–82°N in the years 2006–2009 in ROMS and FESOM.

Eddy type	Radius (km)	Abs. rel. vorticity (normalised by f)	Lifetime (days)	Travel distance (km)
<i>ROMS</i>				
All eddies	4.9 (2.8)	0.40 (0.22)	10 (14)	34 (44)
Cyclones	4.6 (2.6)	0.41 (0.25)	10 (12)	33 (38)
Anticyclones	5.2 (3.0)	0.39 (0.18)	10 (16)	35 (51)
<i>FESOM</i>				
All eddies	5.6 (3.3)	0.28 (0.18)	11 (16)	35 (44)
Cyclones	5.3 (3.1)	0.29 (0.19)	11 (15)	35 (43)
Anticyclones	6.0 (3.4)	0.27 (0.16)	11 (16)	35 (45)




Figure 1. Bathymetry of the Fram Strait. Coloured polygons indicate regions used for analysis: Svalbard shelf (green), West Spitsbergen Current (cyan), central southern Fram Strait (yellow), central northern Fram Strait (blue), and Yermak and Svalbard Branch (magenta). Coloured dots indicate moorings deployed across Fram Strait at 78°50'N; red and magenta dots show moorings used to compute velocity time series representative for the EGC and WSC, respectively. Black arrows show major currents in the Fram Strait (WSC: West Spitsbergen Current, EGC: East Greenland Current, YB: Yermak Branch, SB: Svalbard Branch). MD and YP indicate the locations of the Molloy Deep and the Yermak Plateau, respectively.




Figure 2. Simulated relative vorticity at 100 m depth on Jan 1st, 2006 for ROMS (a) and FESOM (b). Simulated Okubo-Weiss parameter (s^{-2}) at 100 m depth on Jan 1st, 2006 in a region west of Svalbard (grey box in the top panel) for ROMS (c) and FESOM (d). Shown are only values with $OW < -0.2\sigma_{OW}$, where σ_{OW} is the spatial standard deviation of OW at that day. Red arrows show the velocity, with only every 8th vector plotted. Cyan and magenta contours show anticyclonic and cyclonic eddies respectively identified by the Nencioli algorithm (Nencioli et al., 2010). Grey contour lines indicate bathymetry at 1000 m intervals.




Figure 3. Temperature (a,b) and salinity (c,d) in 100 m depth averaged over the time period 2006–2009 simulated by ROMS (left) and FESOM (right). Dots show mooring measurements in 75 m depth for the same time period (von Appen et al., 2019). Grey contour lines indicate bathymetry at 1000 m intervals.




Figure 4. Power density spectra of daily averaged absolute velocity in 75 m depth in the (top) West Spitsbergen Current and (bottom) East Greenland Current from mooring measurements (blue), and models FESOM (red) and ROMS (yellow). Thick lines indicate slopes of the spectra.




Figure 5. Seasonal cycle of eddy kinetic energy at 75 m depth across Fram Strait at 78° 50'N from (a) mooring measurements (von Appen et al., 2019), (b) FESOM, and (c) ROMS for the years 2006–2009.




Figure 6. (a,b) Total number of eddy occurrences in the years 2006–2009 for (left) ROMS and (right) FESOM, binned in a $1/24^\circ$ grid and smoothed with a 3 point Hanning window kernel. (c,d) Difference between cyclonic and anticyclonic eddies (cyclones minus anticyclones).




Figure 7. Histogram of (a) radius, (b) maximum relative vorticity normalised by f , (c) water depth, (d) eddy lifetime and (e) travel distance for anticyclonic (blue) and cyclonic eddies (red) normalised by the number of eddies/tracks tracked in the area 8°W – $20^{\circ}\text{E}/76^{\circ}\text{N}$ – 82°N in the years 2006–2009 in ROMS (dark colours) and FESOM (light colours).




Figure 8. Eddy tracks of cyclones (red lines, a) and anticyclones (blue lines, b) with lifetimes of more than 30 days that are generated in five different regions indicated by coloured polygons, see Fig. 1, detected in simulation ROMS from 2006–2009. Light and dark colours of the lines indicate the beginning and end of the track, respectively.




Figure 9. The same as Figure 8, but for FESOM.




Figure 10. Vertical extent of eddies tracked in 100 m depth during the years 2006–2009 in ROMS (dark colours) and FESOM (light colours) with lifetimes >30 days for cyclones (red) and anticyclones (blue) generated in regions a) Svalbard shelf, b) West Spitsbergen Current, c) southern central Fram Strait, d) northern central Fram Strait and e) Yermak and Svalbard Branch. Left, middle and right panels show relative vorticity, temperature anomaly and salinity anomaly, respectively. Anomalies are calculated by taking the value in the eddy centre relative to the mean value of the month.




Figure 11. Simulated depth integrated energy transfer from (a,b) mean kinetic to eddy kinetic energy (HRS, product of horizontal Reynolds stress and mean shear, $\int_H \left(-\overline{\mathbf{u}'\mathbf{u}'} \cdot \frac{\partial \overline{\mathbf{u}}}{\partial x} - \overline{\mathbf{v}'\mathbf{v}'} \cdot \frac{\partial \overline{\mathbf{u}}}{\partial y} \right) dz$) and (c,d) eddy available potential to eddy kinetic energy (vertical eddy buoyancy flux, $\int_H \overline{w'b'} dz$) averaged for 2006–2009 in (a,c) ROMS and (b,d) FESOM. Black contours show bathymetry contours at 1000 m intervals and at 200 m and 500 m depth.




Figure 12. a) Topographic $\beta = -\frac{f}{H}\nabla H$ and b) $\beta - \partial_{xx}v$, where v is the simulated meridional velocity from FESOM. The second derivative of v is computed from monthly means, and then averaged over the years 2006–2009. A change of sign of $\beta - \partial_{xx}v$ is a necessary condition for barotropic instability. Black contours show bathymetry contours at 1000 m intervals and at 250 m and 500 m depth.




Figure 13. Ertel potential vorticity (left panel) and its gradient in zonal direction (right panel) across Fram Strait at (a) 78°N and (b) 78°50'N computed from long-term mean FESOM data (2006–2009). Black lines show simulated meridional velocity contours (0.1 and 0.2 m/s), and white lines show the simulated 27.9, 28, 28.1, 28.2, and 28.22 kg/m³ isopycnals.